

ON SPECTRAL ESTIMATES FOR TWO-DIMENSIONAL SCHRÖDINGER OPERATORS

A. LAPTEV AND M. SOLOMYAK

ABSTRACT. For a two-dimensional Schrödinger operator $\mathbf{H}_{\alpha V} = -\Delta - \alpha V$, $V \geq 0$, we study the behavior of the number $N_-(\mathbf{H}_{\alpha V})$ of its negative eigenvalues (bound states), as the coupling parameter α tends to infinity. A wide class of potentials is described, for which $N_-(\mathbf{H}_{\alpha V})$ has the semi-classical behavior, i.e., $N_-(\mathbf{H}_{\alpha V}) = O(\alpha)$. For the potentials from this class, the necessary and sufficient condition is found for the validity of the Weyl asymptotic law.

1. INTRODUCTION

1.1. **Preliminaries.** Let $\mathbf{H}_{\alpha V}$ be a Schrödinger operator

$$(1.1) \quad \mathbf{H}_{\alpha V} = -\Delta - \alpha V$$

on \mathbb{R}^2 . We suppose that $V \geq 0$, and $\alpha > 0$ is the coupling constant. We write $N_-(\mathbf{H}_{\alpha V})$ for the number of negative eigenvalues of $\mathbf{H}_{\alpha V}$, counted with multiplicities:

$$N_-(\mathbf{H}_{\alpha V}) = \#\{j \in \mathbb{N} : \lambda_j(\mathbf{H}_{\alpha V}) < 0\}.$$

As it is well known, the lowest possible (semi-classical) rate of growth of this function is

$$(1.2) \quad N_-(\mathbf{H}_{\alpha V}) = O(\alpha), \quad \alpha \rightarrow \infty.$$

This agrees with the Weyl-type asymptotic formula

$$(1.3) \quad \lim_{\alpha \rightarrow \infty} \alpha^{-1} N_-(\mathbf{H}_{\alpha V}) = \frac{1}{4\pi} \int_{\mathbb{R}^2} V dx$$

that is satisfied if the potential behaves fine enough.

The exhaustive description of the classes of potentials on \mathbb{R}^2 , such that (1.2) or (1.3) is satisfied, is unknown till now. This is in contrast with the case of dimensions $d > 2$, where the celebrated Cwikel-Lieb-Rozenblum estimate describes the class of potentials, for which both the estimate $N_-(\mathbf{H}_{\alpha V}) = O(\alpha^{d/2})$ and the Weyl asymptotic formula hold true.

In the forthcoming discussion, $\mathcal{P}_{\text{semi}}$ stands for the class of all potentials $V \geq 0$ on \mathbb{R}^2 , such that (1.2) is satisfied, and $\mathcal{P}_{\text{Weyl}}$ stands for the class of all such potentials that the asymptotics (1.3) holds true. It is clear that

$$(1.4) \quad \mathcal{P}_{\text{Weyl}} \subset \mathcal{P}_{\text{semi}}.$$

The first results describing wide classes of potentials $V \in \mathcal{P}_{\text{semi}}$ were obtained in [9] and [1]. In the latter paper, this was done also for the class $\mathcal{P}_{\text{Weyl}}$. In particular, it was shown there that the inclusion in (1.4) is proper. What is more, in [1] the general nature of potentials $V \in \mathcal{P}_{\text{semi}} \setminus \mathcal{P}_{\text{Weyl}}$ was explained.

1991 *Mathematics Subject Classification.* 35J10; 35P20.

Key words and phrases. Schrödinger operator on \mathbb{R}^2 ; bound states; spectral estimates.

Some further estimates guaranteeing $V \in \mathcal{P}_{\text{semi}}$ were obtained in the recent paper [4]. We would like to mention also the paper [8] whose authors have obtained some new results that give for $N_-(\mathbf{H}_{\alpha V})$ the order of growth larger than $O(\alpha)$.

In the papers [5, 3] the important case of radial potentials, $V(x) = F(|x|)$, was analyzed. For such potentials in [3] an integral estimate for $N_-(\mathbf{H}_{\alpha V})$ was obtained guaranteeing the inclusion $V \in \mathcal{P}_{\text{semi}}$ (actually, it guarantees also that $V \in \mathcal{P}_{\text{Weyl}}$). This result was strengthened in the recent paper [7] where for the radial potentials the necessary and sufficient conditions for $V \in \mathcal{P}_{\text{semi}}$ and for $V \in \mathcal{P}_{\text{Weyl}}$ were established.

In the present paper we return to the study of general (that is, not necessarily radial) potentials. We obtain an estimate that covers the main results of [1, 4]. It does not cover the estimate obtained in [9], however it has an important advantage compared with the latter: it does not use the intricate Orlicz norms appearing in [9].

In the papers [5, 3] the important case of radial potentials, $V(x) = F(|x|)$, was analyzed. For such potentials in [3] an integral estimate for $N_-(\mathbf{H}_{\alpha V})$ was obtained guaranteeing the inclusion $V \in \mathcal{P}_{\text{semi}}$. Actually, it guarantees also that $V \in \mathcal{P}_{\text{Weyl}}$. This result was strengthened in the recent paper [7] where for the radial potentials the necessary and sufficient conditions for $V \in \mathcal{P}_{\text{semi}}$ and for $V \in \mathcal{P}_{\text{Weyl}}$ were established.

In the present paper we return to the study of general (that is, not necessarily radial) potentials. We obtain an estimate that covers the main results of [1]. It does not cover the estimate obtained in [9], however it has an important advantage compared with the latter: it does not use the intricate Orlicz norms appearing in [9].

1.2. Formulation of the main result. Below (r, ϑ) stand for the polar coordinates in \mathbb{R}^2 , and \mathbb{S} stands for the unit circle $r = 1$. Given a function V , such that $V(r, \cdot) \in L_1(\mathbb{S})$ for almost all $r > 0$, we introduce its radial and non-radial parts

$$V_{\text{rad}}(r) = \frac{1}{2\pi} \int_{\mathbb{S}} V(r, \vartheta) d\vartheta; \quad V_{\text{nrad}}(r, \vartheta) = V(r, \vartheta) - V_{\text{rad}}(r).$$

In our result the conditions will be imposed separately on the radial and on the non-radial parts of a given potential V . For handling the radial part, we need some auxiliary operator family on the real line, of the form

$$(1.5) \quad (\mathbf{M}_{\alpha G}\varphi)(t) = -\varphi''(t) - \alpha G(t)\varphi(t), \quad \varphi(0) = 0,$$

with the "effective potential"

$$(1.6) \quad G(t) = G_V(t) = e^{2|t|} V_{\text{rad}}(e^t).$$

Due to the condition $\varphi(0) = 0$ in (1.5), for every α the operator $\mathbf{M}_{\alpha G}$ is the direct sum of two operators, each acting on the half-line. The sharp spectral estimates for $\mathbf{M}_{\alpha G}$ can be given in terms of the number sequence (see Eq. (1.13) in [7])

$$(1.7) \quad \widehat{\mathfrak{z}}(G) = \{\widehat{\zeta}_j(G)\}_{j \geq 0} : \quad \widehat{\zeta}_0(G) = \int_{D_0} G(t) dt, \quad \widehat{\zeta}_j(G) = \int_{|t| \in D_j} |t| G(t) dt \quad (j \in \mathbb{N})$$

where $D_0 = (-1, 1)$ and $D_j = (e^{j-1}, e^j)$ for $j \in \mathbb{N}$. For our purposes, it is convenient to express properties of this sequence in terms of the "weak ℓ_q -spaces" $\ell_{q,\infty}$. Actually, in the main body of this paper we deal only with $q = 1$, and below we remind the definition of $\ell_{1,\infty}$. The definition of the weak ℓ_q -spaces with $q \neq 1$ can be found, e.g., in [1], Subsection 1.4.

Given a sequence of real numbers $\mathbf{x} = \{x_j\}_{j \in \mathbb{N}}$, such that $x_j \rightarrow 0$, we denote

$$n_+(\varepsilon, \mathbf{x}) = \#\{j : |x_j| > \varepsilon\}, \quad \varepsilon > 0.$$

The sequence \mathbf{x} belongs to $\ell_{1,\infty}$, if

$$\|\mathbf{x}\|_{1,\infty} := \sup_{\varepsilon > 0} (\varepsilon n_+(\varepsilon, \mathbf{x})) < \infty.$$

This is a linear space, and the functional $\|\cdot\|_{1,\infty}$ defines a quasinorm in it. The latter means that, instead of the standard triangle inequality, this functional meets a weaker property:

$$\|\mathbf{x} + \mathbf{y}\|_{1,\infty} \leq c(\|\mathbf{x}\|_{1,\infty} + \|\mathbf{y}\|_{1,\infty}),$$

with some constant $c > 1$ that does not depend on the sequences \mathbf{x}, \mathbf{y} . This quasinorm defines a topology in $\ell_{1,\infty}$; there is no norm compatible with this topology.

The space $\ell_{1,\infty}$ is non-separable. Consider its closed subspace $\ell_{1,\infty}^\circ$ in which the sequences \mathbf{x} with only a finitely many non-zero terms form a dense subset. This subspace is separable, and its elements are characterized by the property

$$\mathbf{x} \in \ell_{1,\infty}^\circ \iff \varepsilon n_+(\varepsilon, \mathbf{x}) \rightarrow 0, \quad \varepsilon \rightarrow 0.$$

The (non-linear) functionals

$$(1.8) \quad \Delta_1(\mathbf{x}) = \limsup_{\varepsilon \rightarrow 0} (\varepsilon n_+(\varepsilon, \mathbf{x})), \quad \delta_1(\mathbf{x}) = \liminf_{\varepsilon \rightarrow 0} (\varepsilon n_+(\varepsilon, \mathbf{x}))$$

are well-defined on the space $\ell_{1,\infty}$, and

$$\delta_1(\mathbf{x}) \leq \Delta_1(\mathbf{x}) \leq \|\mathbf{x}\|_{1,\infty}.$$

It is clear that $\ell_{1,\infty}^\circ = \{\mathbf{x} \in \ell_{1,\infty} : \Delta_1(\mathbf{x}) = 0\}$.

The conditions on V_{nrad} will be given in terms of the space $L_1(\mathbb{R}_+, L_p(\mathbb{S}))$, with an arbitrarily chosen $p > 1$. This is the function space on \mathbb{R}^2 , with the following norm:

$$(1.9) \quad \|f\|_{L_1(\mathbb{R}_+, L_p(\mathbb{S}))} = \int_{\mathbb{R}_+} \left(\int_{\mathbb{S}} |f(r, \vartheta)|^p d\vartheta \right)^{1/p} r dr.$$

This is a separable Banach space, and the bounded functions whose support is a compact subset in $\mathbb{R}^2 \setminus \{0\}$ are dense in it. The space $L_1(\mathbb{R}_+, L_p(\mathbb{S}))$ was used in the paper [6], and its results are one of the basic tools in our proof below.

Here is the main result of the paper.

Theorem 1.1. *Let a potential $V \geq 0$ be such that $\widehat{\mathfrak{z}}(G_V) \in \ell_{1,\infty}$, and*

$$(1.10) \quad V_{\text{nrad}} \in L_1(\mathbb{R}_+, L_p(\mathbb{S})) \quad \text{with some } p > 1.$$

Then $V \in \mathcal{P}_{\text{semi}}$, and the estimate is satisfied

$$(1.11) \quad N_-(\mathbf{H}_{\alpha V}) \leq 1 + C(p)\alpha \left(\|V_{\text{nrad}}\|_{L_1(\mathbb{R}_+, L_p(\mathbb{S}))} + \|\widehat{\mathfrak{z}}(G_V)\|_{\ell_{1,\infty}} \right).$$

Moreover, the following equalities hold true:

$$(1.12) \quad \begin{cases} \limsup_{\alpha \rightarrow \infty} \alpha^{-1} N_-(\mathbf{H}_{\alpha V}) = \frac{1}{4\pi} \int V dx + \limsup_{\alpha \rightarrow \infty} \alpha^{-1} N_-(\mathbf{M}_{\alpha G_V}), \\ \liminf_{\alpha \rightarrow \infty} \alpha^{-1} N_-(\mathbf{H}_{\alpha V}) = \frac{1}{4\pi} \int V dx + \liminf_{\alpha \rightarrow \infty} \alpha^{-1} N_-(\mathbf{M}_{\alpha G_V}). \end{cases}$$

In particular, under the assumption (1.10) the condition $\widehat{\mathfrak{z}}(G_V) \in \ell_{1,\infty}^\circ$ is necessary and sufficient for $V \in \mathcal{P}_{\text{Weyl}}$.

In (1.12), and later on, the integral with no domain specified always means $\int_{\mathbb{R}^2}$.

The formula (1.12), and especially, its proof in Subsection 3.3, show that, in a certain sense, the parts V_{rad} and V_{nrad} contribute to the asymptotic behavior of $N_-(\mathbf{H}_{\alpha V})$ independently. It may also happen that the contribution of V_{rad} is stronger than that of V_{nrad} , and "screens" the latter. This situation is described by the following statement, that complements our main theorem.

Proposition 1.2. *Let a potential $V \geq 0$ be such that $\widehat{\mathfrak{J}}(G_V) \in \ell_{q,\infty}$ with some $q > 1$, and (1.10) is satisfied. Then*

$$\begin{cases} \limsup_{\alpha \rightarrow \infty} \alpha^{-q} N_-(\mathbf{H}_{\alpha V}) = \limsup_{\alpha \rightarrow \infty} \alpha^{-q} N_-(\mathbf{M}_{\alpha G_V}), \\ \liminf_{\alpha \rightarrow \infty} \alpha^{-q} N_-(\mathbf{H}_{\alpha V}) = \liminf_{\alpha \rightarrow \infty} \alpha^{-q} N_-(\mathbf{M}_{\alpha G_V}). \end{cases}$$

This is an analog of statement (b) in Theorem 5.1 of the paper [1]. Its proof is basically the same, and we do not reproduce it here. In the same paper one finds also examples that illustrate the situation described by Proposition 1.2.

2. AUXILIARY MATERIAL

The proof of Theorem 1.1 mainly follows the line worked out in [1] and [9]. The same approach was used in [7], and the material below, in part, duplicates the contents of its Section 2. We systematically use the variational description of the spectrum. In particular, we often define a self-adjoint operator via its corresponding Rayleigh quotient.

2.1. Classes Σ_1, Σ_1° of compact operators. If \mathbf{T} is a linear compact operator in a Hilbert space, then, as usual, $\{s_j(\mathbf{T})\}$ stands for the sequence of its singular numbers, i.e., for the eigenvalues of the non-negative, self-adjoint operator $(\mathbf{T}^* \mathbf{T})^{1/2}$. By $n_+(\varepsilon, \mathbf{T})$ we denote the distribution function of the singular numbers,

$$n_+(\varepsilon, \mathbf{T}) = \#\{j : s_j > \varepsilon\}, \quad \varepsilon > 0.$$

We say that \mathbf{T} belongs to the class Σ_1 if and only if $\{s_j(\mathbf{T})\} \in \ell_{1,\infty}$, and to the class Σ_1° if and only if $\{s_j(\mathbf{T})\} \in \ell_{1,\infty}^\circ$. These are linear, quasinormed spaces with respect to the quasinorm $\|\mathbf{T}\|_{1,\infty}$ induced by this definition. The space Σ_1 is non-separable, and Σ_1° is its separable subspace in which the finite rank operators form a dense subset. Similarly to (1.8), we define the functionals

$$\Delta_1(\mathbf{T}) = \Delta_1(\{s_j(\mathbf{T})\}), \quad \delta_1(\mathbf{T}) = \delta_1(\{s_j(\mathbf{T})\}).$$

Note that

$$\delta_1(\mathbf{T}) \leq \Delta_1(\mathbf{T}) \leq \|\mathbf{T}\|_{1,\infty}.$$

See [2], Section 11.6 for more detail about these spaces, and about the similar spaces Σ_q, Σ_q° for any $q > 0$.

2.2. Reduction of the main problem to compact operators. Let us introduce two subspaces in $C_0^\infty(\mathbb{R}^2)$:

$$\mathcal{F}_0 = \{f \in C_0^\infty : f(x) = \varphi(r), \varphi(1) = 0\}; \quad \mathcal{F}_1 = \{f \in C_0^\infty : \int_0^{2\pi} f(r, \vartheta) d\vartheta = 0, \forall r > 0\}.$$

They are orthogonal to each other both in the L_2 -metric and in the metric of the Dirichlet integral. The Hardy inequalities have a different form on \mathcal{F}_0 and on \mathcal{F}_1 :

$$(2.1) \quad \int \frac{|f(x)|^2}{|x|^2 \ln^2 |x|} dx \leq \frac{1}{4} \int |\nabla f(x)|^2 dx, \quad f \in \mathcal{F}_0;$$

$$(2.2) \quad \int \frac{|f(x)|^2}{|x|^2} dx \leq \int |\nabla f(x)|^2 dx, \quad f \in \mathcal{F}_1.$$

For proving (2.1), one substitutes $r = |x| = e^t$, and then applies the standard Hardy inequality in dimension 1. The proof of (2.2) is quite elementary, it can be found, e.g., in [9], or in [1].

Let us consider the completions $\mathcal{H}_0^1, \mathcal{H}_1^1$ of the spaces $\mathcal{F}_0, \mathcal{F}_1$ in the metric of the Dirichlet integral. It follows from the Hardy inequalities (2.1), (2.2) that these are Hilbert function spaces, embedded into the weighted L_2 , with the weights defined by these inequalities. Consider also their orthogonal sum

$$(2.3) \quad \mathcal{H}^1 = \mathcal{H}_0^1 \oplus \mathcal{H}_1^1.$$

An independent definition of this Hilbert space is

$$\mathcal{H}^1 = \{f \in H_{\text{loc}}^1(\mathbb{R}^2) : \int_0^{2\pi} f(1, \vartheta) d\vartheta = 0, |\nabla f| \in L_2(\mathbb{R}^2)\},$$

with the metric of the Dirichlet integral.

We also define the spaces H_0^1, H_1^1 which are the completions of $\mathcal{F}_0, \mathcal{F}_1$ in $H^1(\mathbb{R}^2)$, and

$$\tilde{H}^1 = H_0^1 \oplus H_1^1 = \{f \in H^1(\mathbb{R}^2) : \int_0^{2\pi} f(1, \vartheta) d\vartheta = 0\}.$$

This is a subspace in $H^1(\mathbb{R}^2)$ of codimension 1.

Finally, we need the spaces $\mathcal{G}_0, \mathcal{G}_1$ which are the completions of $\mathcal{F}_0, \mathcal{F}_1$ in the L_2 -metric. Note that the condition $\varphi(1) = 0$, occuring in the description of \mathcal{F}_0 , disappears for general $f \in \mathcal{G}_0$.

Suppose that $V \geq 0$ is a measurable function, such that

$$(2.4) \quad \mathbf{b}_V[u] := \int V|u|^2 dx \leq C \int |\nabla u|^2 dx, \quad \forall u \in \mathcal{H}^1.$$

Under the assumption (2.4) the quadratic form \mathbf{b}_V defines a bounded self-adjoint operator $\mathbf{B}_V \geq 0$ in \mathcal{H}^1 . If (and only if) this operator is compact, then, by the Birman-Schwinger principle, the quadratic form

$$(2.5) \quad \int (|\nabla u|^2 - \alpha V|u|^2) dx$$

with the form-domain \tilde{H}^1 is closed and bounded from below for each $\alpha > 0$, the negative spectrum of the associated self-adjoint operator $\tilde{\mathbf{H}}_{\alpha V}$ on $L_2(\mathbb{R}^2)$ is finite, and the following equality for the number of its negative eigenvalues holds true:

$$(2.6) \quad N_-(\tilde{\mathbf{H}}_{\alpha V}) = n_+(\alpha^{-1}, \mathbf{B}_V), \quad \forall \alpha > 0.$$

Now, let us withdraw the rank one condition $\int_0^{2\pi} u(1, \vartheta) d\vartheta = 0$ from the description of the form-domain. Then the resulting quadratic form corresponds to the Schrödinger operator $\mathbf{H}_{\alpha V}$ as in (1.1). Hence,

$$N_-(\tilde{\mathbf{H}}_{\alpha V}) \leq N_-(\mathbf{H}_{\alpha V}) \leq N_-(\tilde{\mathbf{H}}_{\alpha V}) + 1,$$

and, by (2.6),

$$n_+(\alpha^{-1}, \mathbf{B}_V) \leq N_-(\mathbf{H}_{\alpha V}) \leq n_+(\alpha^{-1}, \mathbf{B}_V) + 1.$$

Thus, the study of the quantity $N_-(\mathbf{H}_{\alpha V})$ for all $\alpha > 0$ is reduced to the investigation of the "individual" operator \mathbf{B}_V , which is nothing but the Birman-Schwinger operator for the family of operators in $L_2(\mathbb{R}^2)$ associated with the family of quadratic forms in (2.5). Note that the Birman-Schwinger operator for the original family (1.1) is ill-defined, since the completion of the space $H^1(\mathbb{R}^2)$ in the metric of the Dirichlet integral is not a space of functions on \mathbb{R}^2 .

3. PROOF OF THEOREM 1.1

3.1. Decomposition of the quadratic form \mathbf{b}_V . Given a function $u \in \mathcal{H}^1$, we agree to standardly denote its components in the decomposition (2.3) by $\varphi(r), v(r, \vartheta)$. Along with the quadratic form \mathbf{b}_V , we consider its "parts" in the subspaces $\mathcal{H}_0^1, \mathcal{H}_1^1$:

$$\mathbf{b}_{V,0}[u] = \mathbf{b}_V[\varphi], \quad \mathbf{b}_{V,1}[u] = \mathbf{b}_V[v].$$

Let $\mathbf{B}_{V,j}$, $j = 0, 1$, stand for the corresponding self-adjoint operators in \mathcal{H}_j^1 . Using the orthogonal decomposition (2.3), we see that

$$(3.1) \quad \mathbf{b}_V[u] = \mathbf{b}_{V,0}[\varphi] + \mathbf{b}_{V,1}[v] + 2 \int V(x) \operatorname{Re}(\varphi(|x|) \overline{v(x)}) dx.$$

For the radial potentials the last term vanishes, and this considerably simplifies the reasoning, see [7]. For the general potentials this is no more true. Still, the following inequality is always valid:

$$(3.2) \quad \mathbf{b}_V[u] \leq 2(\mathbf{b}_V[\varphi] + \mathbf{b}_V[v]),$$

and it shows that for estimation of $\|\mathbf{B}_V\|_{1,\infty}$ it suffices to evaluate the quasinorms in Σ_1 of the operators $\mathbf{B}_{V,0}, \mathbf{B}_{V,1}$ separately.

The estimation of $\|\mathbf{B}_{V,0}\|_{1,\infty}$ will be based upon the following result on the operators \mathbf{F}_G on real line, whose Rayleigh quotient is

$$(3.3) \quad \frac{\int_{\mathbb{R}} G(t) |\omega(t)|^2 dt}{\int_{\mathbb{R}} |\omega'(t)|^2 dt}, \quad \omega(0) = 0.$$

Clearly, this is the Birman-Schwinger operator for the family $\mathbf{M}_{\alpha G}$ given by (1.5).

Proposition 3.1. *Let a function $G \in L_{1,loc}(\mathbb{R})$, $G \geq 0$, be given. Define the corresponding number sequence $\widehat{\mathfrak{z}}(G)$ as in (1.7), and suppose that $\widehat{\mathfrak{z}}(G) \in \ell_{1,\infty}$. Then the operator \mathbf{F}_G is well-defined, belongs to the class Σ_1 , and the estimate is satisfied,*

$$(3.4) \quad \|\mathbf{F}_G\|_{1,\infty} \leq C \|\widehat{\mathfrak{z}}(G)\|_{1,\infty}.$$

If $\widehat{\mathfrak{z}}(G) \in \ell_{1,\infty}^\circ$, then $\widehat{\mathfrak{z}}(G) \in \Sigma_1^\circ$.

For the proof, see Section 4 in the paper [1]. There the operators on the half-line were considered, however the passage to the case of the whole line is straightforward, due to the condition $\omega(0) = 0$ in (3.3). In this respect, see also a discussion in [7], Section 3.

Now we turn to the operator $\mathbf{B}_{V,1}$. The estimation of its quasinorm in Σ_1 uses a result that is a particular case (for $l = 1$) of Theorem 1.2 in the paper [6]. We present its equivalent formulation, more convenient for our purposes. Namely, we formulate it for the Birman-Schwinger operator, rather than for the original Schrödinger operator, as it was done in [6].

Proposition 3.2. *Let $V \geq 0$, $V \in L_1(\mathbb{R}_+, L_p(\mathbb{S}))$, with some $p > 1$. Then the operator $\widehat{\mathbf{B}}_V$, whose Rayleigh quotient is*

$$(3.5) \quad \frac{\int V(x)|u|^2 dx}{\int (|\nabla u|^2 + |x|^{-2}|u|^2) dx}, \quad u \in \mathcal{H}_1^1,$$

belongs to the class Σ_1 , and

$$(3.6) \quad \|\widehat{\mathbf{B}}_V\|_{1,\infty} \leq C(p)\|V\|_{L_1(\mathbb{R}_+, L_p(\mathbb{S}))}.$$

We recall that the norm appearing in (3.6) was defined in (1.9).

3.2. Proof of (1.11). As it was explained in the previous subsection, we have to estimate the quasinorms of the operators $\mathbf{B}_{V,0}$, $\mathbf{B}_{V,1}$ in the space Σ_1 .

Consider first the operator $\mathbf{B}_{V,0}$. The corresponding Rayleigh quotient is

$$(3.7) \quad \frac{\int_{\mathbb{R}^2} V(r, \vartheta)|\varphi(r)|^2 r dr d\vartheta}{\int_{\mathbb{R}^2} |\varphi'(r)|^2 r dr d\vartheta} = \frac{\int_0^\infty V_{\text{rad}}(r)|\varphi(r)|^2 r dr}{\int_0^\infty |\varphi'(r)|^2 r dr}.$$

The standard substitution $r = e^t$, $\varphi(r) = \omega(t)$; $t \in \mathbb{R}$, reduces it to the form

$$\frac{\int_{\mathbb{R}} G_V(t)|\omega(t)|^2 dt}{\int_{\mathbb{R}} |\omega'(t)|^2 dt}, \quad \omega(0) = 0.$$

where the potential G_V is given by (1.6). Now, Proposition 3.1 applies, and we arrive at the estimate

$$\|\mathbf{B}_{V,0}\|_{1,\infty} \leq C\|\widehat{\mathfrak{F}}(G)\|_{1,\infty}.$$

The Rayleigh quotient for the operator $\mathbf{B}_{V,1}$ is given by

$$\frac{\int V(x)|u|^2 dx}{\int |\nabla u|^2 dx}, \quad u \in \mathcal{H}_1^1.$$

Due to the Hardy inequality (2.2), on the subspace \mathcal{H}_1^1 the norm of the Dirichlet integral is equivalent to the norm generated by the quadratic form in the denominator of (3.5). Hence, the estimate (3.6) applies to this operator, with some other constant factor $C'(p)$. So, we have

$$(3.8) \quad \|\mathbf{B}_{V,1}\|_{1,\infty} \leq C'(p)\|V\|_{L_1(\mathbb{R}_+, L_p(\mathbb{S}))}.$$

The estimates (3.4) and (3.8), together with the inequality (3.2), imply the desired estimate (1.11).

3.3. Proof of (1.12). First of all, we are going to show that

$$(3.9) \quad \lim_{\varepsilon \rightarrow 0} (\varepsilon n_+(\varepsilon, \mathbf{B}_{V,1})) = \frac{1}{4\pi} \int_{\mathbb{R}^2} V dx.$$

For $V \in C_0^\infty(\mathbb{R}^2 \setminus \{0\})$, Theorem 5.1 in [1] yields that $N_-(\mathbf{H}_{\alpha V}) \sim (4\pi)^{-1} \alpha \int V dx$ as $\alpha \rightarrow \infty$. By the Birman-Schwinger principle, this is equivalent to $n_+(\varepsilon, \mathbf{B}_V) \sim (4\pi\varepsilon)^{-1} \int V dx$ as $\varepsilon \rightarrow 0$. The spectrum of $\mathbf{B}_{V,1}$ has the same asymptotic behavior, since for such potentials the subspace \mathcal{H}_0^1 does not contribute to the asymptotic coefficient.

Now, let $V \geq 0$ be an arbitrary potential from $L_1(\mathbb{R}_+, L_p(\mathbb{S}))$. Then, approximating it by the functions from C_0^∞ and taking into account the continuity of the asymptotic coefficients in the metric of Σ_1 (see [2], Theorem 11.6.6), we extend the formula to all such V . So, (3.9) is established.

Return to the study of the operator \mathbf{B}_V . Along with it, let us consider the direct orthogonal sum $\mathcal{B}_V = \mathbf{B}_{V,0} \oplus \mathbf{B}_{V,1}$. Evidently,

$$n_+(\varepsilon, \mathcal{B}_V) = n_+(\varepsilon, \mathbf{B}_{V,0}) + n_+(\varepsilon, \mathbf{B}_{V,1}).$$

Hence, for justifying the asymptotic formulas (1.12) it suffices to show that the off-diagonal term in (3.1) generates an operator of the class Σ_1° . To this end, we first of all note that

$$(3.10) \quad \int V \operatorname{Re}(\varphi \bar{v}) dx = \int V_{\text{nrad}} \operatorname{Re}(\varphi \bar{v}) dx,$$

since v is orthogonal (in L_2) to all functions depending only on $|x|$.

Suppose now that the function V_{nrad} has a compact support in $\mathbb{R}^2 \setminus \{0\}$. Then the integral in the right-hand side of (3.10) is actually taken over some annulus $a \leq r \leq a^{-1}$, $a < 1$. Hence,

$$\begin{aligned} 2 \left| \int V_{\text{nrad}} \operatorname{Re}(\varphi \bar{v}) dx \right| &\leq \delta \int_a^{a^{-1}} r dr \int_{\mathbb{S}} |V_{\text{nrad}}(r, \vartheta)| |v(r, \vartheta)|^2 d\vartheta \\ &+ \delta^{-1} \int_a^{a^{-1}} r dr \int_{\mathbb{S}} |V_{\text{nrad}}(r, \vartheta)| |\varphi(r)|^2 d\vartheta. \end{aligned}$$

The first term on the right generates an operator on \mathcal{H}_1^1 , say, \mathbf{T}_1 , to which the estimate (3.8) applies, and it gives

$$\|\mathbf{T}_1\|_{1,\infty} \leq C'(p)\delta.$$

The second term generates an operator on \mathcal{H}_0^1 , say, \mathbf{T}_0 . Its Rayleigh quotient is of the type (3.7) but with the integration over a compact subset in $(0, \infty)$. It follows that the spectrum of \mathbf{T}_0 obeys Weyl's asymptotic law, $\lambda_j(\mathbf{T}_0) \asymp c_j^{-2}$, and hence, $\mathbf{T}_0 \in \Sigma_1^\circ$. Taking δ arbitrarily small, we conclude that the asymptotics (1.12) is satisfied in the case where V_{nrad} is compactly supported.

Finally, we approximate the function V_{nrad} by compactly supported functions in the metric (1.9), and again apply Theorem 11.6.5 from the book [2]. This extends the asymptotic formula (1.12) to all potentials, that meet the conditions of Theorem 1.1, and thus, concludes the proof.

REFERENCES

- [1] Birman, M.Sh., Laptev, A.: *The negative discrete spectrum of a two-dimensional Schrödinger operator*, Comm. Pure Appl. Math. **49**, no. 9, 967–997 (1996).
- [2] Birman, M.Sh., Solomyak, M.: *Spectral theory of selfadjoint operators in Hilbert space*, D. Reidel Publishing Co., Dordrecht, 1987.
- [3] Chadan, K., Khuri, N. N., Martin, A., Wu, Tai Tsun: *Bound states in one and two spatial dimensions*, J. Math. Phys. **44**, no. 2, 406–422 (2003).
- [4] Grigor'yan, A., Nadirashvili, N.: *Negative eigenvalues of two-dimensional Schrödinger operators*, arXiv:1112.4986
- [5] Laptev, A.: *The negative spectrum of the class of two-dimensional Schrödinger operators with potentials that depend on the radius*, (Russian) Funktsional. Anal. i Prilozhen. **34**, no. 4, 85–87 (2000); translation in Funct. Anal. Appl. **34**, no. 4, 305–307 (2000).
- [6] Laptev, A., Netrusov, Yu.: *On the negative eigenvalues of a class of Schrödinger operators*, Differential operators and spectral theory, 173–186, Amer. Math. Soc. Transl. Ser. 2, **189**, Amer. Math. Soc., Providence, RI, 1999.
- [7] Laptev, A., Solomyak, M.: *On the negative spectrum of the two-dimensional Schrödinger operator with radial potential*, Comm. Math. Phys., in press.
- [8] Molchanov, S., Vainberg, B.: *On negative eigenvalues of low-dimensional Schrödinger operators*, arXiv:1105.0937.
- [9] Solomyak, M.: *Piecewise-polynomial approximation of functions from $H^l((0,1)^d)$, $2l = d$, and applications to the spectral theory of the Schrödinger operator*, Israel J. Math. **86**, no. 1-3, 253–275 (1994).

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, HUXLEY BUILDING, 180 QUEEN'S GATE, LONDON SW7 2AZ, UK

E-mail address: a.laptev@imperial.ac.uk

DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE, REHOVOT, ISRAEL

E-mail address: michail.solomyak@weizmann.ac.il