

**Lecture 10 2.11.2015**

*Kolmogorov's approach: conditional expectations via  $\sigma$ -fields*

The problem with the approach of L9 (discrete and density cases) is that joint densities need not exist – do not exist, in general. One of the great contributions of Kolmogorov's classic book of 1933 was the realization that measure theory – specifically, the Radon-Nikodym theorem – provides a way to treat conditioning in general, without assuming that we are in the discrete case or density case above.

Recall that the probability triple is  $(\Omega, \mathcal{F}, \mathcal{P})$ . Take  $\mathcal{B}$  a sub- $\sigma$ -field of  $\mathcal{F}$ ,  $\mathcal{B} \subset \mathcal{F}$  (recall: a  $\sigma$ -field represents information; the big  $\sigma$ -field  $\mathcal{F}$  represents 'knowing everything', the small  $\sigma$ -field  $\mathcal{B}$  represents 'knowing something').

Suppose that  $Y$  is a non-negative random variable whose expectation exists:  $E[Y] < \infty$ . The set-function

$$Q(B) := \int_B Y dP \quad (B \in \mathcal{B})$$

is non-negative (because  $Y$  is),  $\sigma$ -additive – because

$$\int_B Y dP = \sum_n \int_{B_n} Y dP$$

if  $B = \cup_n B_n$ ,  $B_n$  disjoint – and defined on the  $\sigma$ -algebra  $\mathcal{B}$ , so is a *measure* on  $\mathcal{B}$ . If  $P(B) = 0$ , then  $Q(B) = 0$  also (the integral of anything over a null set is zero), so  $Q \ll P$ . By the Radon-Nikodym theorem (II.4), there exists a Radon-Nikodym derivative of  $Q$  with respect to  $P$  on  $\mathcal{B}$ , which is  $\mathcal{B}$ -measurable [in the Radon-Nikodym theorem as stated in II.4, we had  $\mathcal{F}$  in place of  $\mathcal{B}$ , and got a random variable, i.e. an  $\mathcal{F}$ -measurable function. Here, we just replace  $\mathcal{F}$  by  $\mathcal{B}$ .] Following Kolmogorov (1933), we call this Radon-Nikodym derivative the *conditional expectation of  $Y$  given (or conditional on)  $\mathcal{B}$* ,  $E[Y|\mathcal{B}]$ : this is  $\mathcal{B}$ -measurable, integrable, and satisfies

$$\int_B Y dP = \int_B E[Y|\mathcal{B}] dP \quad \forall B \in \mathcal{B}. \quad (*)$$

In the general case, where  $Y$  is a random variable whose expectation exists ( $E[|Y|] < \infty$ ) but which can take values of both signs, decompose  $Y$  as

$$Y = Y_+ - Y_-$$

and define  $E(Y|\mathcal{B})$  by linearity as

$$E[Y|\mathcal{B}] := E[Y_+|\mathcal{B}] - E[Y_-|\mathcal{B}].$$

Suppose now that  $\mathcal{B}$  is the  $\sigma$ -field generated by a random variable  $X$ :  $\mathcal{B} = \sigma(X)$  (so  $\mathcal{B}$  represents the information contained in  $X$ , or what we know when we know  $X$ ). Then  $E[Y|\mathcal{B}] = E[Y|\sigma(X)]$ , which is written more simply as  $E[Y|X]$ . Its defining property is

$$\int_B Y dP = \int_B E[Y|X] dP \quad \forall B \in \sigma(X).$$

Similarly, if  $\mathcal{B} = \sigma(X_1, \dots, X_n)$  ( $\mathcal{B}$  is the information in  $(X_1, \dots, X_n)$ ) we write  $E[Y|\sigma(X_1, \dots, X_n)]$  as  $E[Y|X_1, \dots, X_n]$ :

$$\int_B Y dP = \int_B E[Y|X_1, \dots, X_n] dP \quad \forall B \in \sigma(X_1, \dots, X_n).$$

**Note.** 1. To check that something is a conditional expectation: we have to check that it integrates the right way over the right sets [i.e., as in (\*)].

2. From (\*): if two things integrate the same way over all sets  $B \in \mathcal{B}$ , they have the same conditional expectation given  $\mathcal{B}$ .

3. For notational convenience, we use  $E[Y|\mathcal{B}]$  and  $E_{\mathcal{B}}Y$  interchangeably.

4. The conditional expectation thus defined coincides with any we may have already encountered - in regression or multivariate analysis, for example. However, this may not be immediately obvious. The conditional expectation defined above - via  $\sigma$ -fields and the Radon-Nikodym theorem - is rightly called by Williams ([W], p.84) ‘the central definition of modern probability’. It may take a little getting used to. As with all important but non-obvious definitions, it proves its worth in action: see II.6 below for properties of conditional expectations, and Chapter III for stochastic processes, particularly martingales [defined in terms of conditional expectations].

## §6. Properties of Conditional Expectations.

1.  $\mathcal{B} = \{\emptyset, \Omega\}$ . Here  $\mathcal{B}$  is the *smallest* possible  $\sigma$ -field (*any*  $\sigma$ -field of subsets of  $\Omega$  contains  $\emptyset$  and  $\Omega$ ), and represents ‘knowing nothing’.

$$E[Y|\{\emptyset, \Omega\}] = EY.$$

*Proof.* We have to check (\*) of §5 for  $B = \emptyset$  and  $B = \Omega$ . For  $B = \emptyset$  both sides are zero; for  $B = \Omega$  both sides are  $EY$ . //

2.  $\mathcal{B} = \mathcal{F}$ . Here  $\mathcal{B}$  is the *largest* possible  $\sigma$ -field: ‘knowing everything’.

$$E[Y|\mathcal{F}] = Y \quad P - a.s.$$

*Proof.* We have to check (\*) for *all* sets  $B \in \mathcal{F}$ . The only integrand that integrates like  $Y$  over *all* sets is  $Y$  itself, or a function agreeing with  $Y$  except on a set of measure zero.

*Note.* When we condition on  $\mathcal{F}$  (‘knowing everything’), we *know*  $Y$  (because we know everything). There is thus no uncertainty left in  $Y$  to average out, so taking the conditional expectation (averaging out remaining randomness) has no effect, and leaves  $Y$  unaltered.

3. If  $Y$  is  $\mathcal{B}$ -measurable,  $E[Y|\mathcal{B}] = Y \quad P - a.s.$

*Proof.* Recall that  $Y$  is *always*  $\mathcal{F}$ -measurable (this is the definition of  $Y$  being a random variable). For  $\mathcal{B} \subset \mathcal{F}$ ,  $Y$  may not be  $\mathcal{B}$ -measurable, but if it is, the proof above applies with  $\mathcal{B}$  in place of  $\mathcal{F}$ .

*Note.* If  $Y$  is  $\mathcal{B}$ -measurable, when we are given  $\mathcal{B}$  (that is, when we condition on it), we *know*  $Y$ . That makes  $Y$  effectively a constant, and when we take the expectation of a constant, we get the same constant.

4. If  $Y$  is  $\mathcal{B}$ -measurable,  $E[YZ|\mathcal{B}] = YE[Z|\mathcal{B}] \quad P - a.s.$

We refer for the proof of this to [W], p.90, proof of (j).

*Note.* Williams calls this property ‘taking out what is known’. To remember it: if  $Y$  is  $\mathcal{B}$ -measurable, then given  $\mathcal{B}$  we know  $Y$ , so  $Y$  is effectively a constant, so can be taken out through the integration signs in (\*), which is what we have to check (with  $YZ$  in place of  $Y$ ).

5. If  $\mathcal{C} \subset \mathcal{B}$ ,  $E[E[Y|\mathcal{B}]|\mathcal{C}] = E[Y|\mathcal{C}] \quad a.s.$

*Proof.*  $E_{\mathcal{C}}E_{\mathcal{B}}Y$  is  $\mathcal{C}$ -measurable, and for  $C \in \mathcal{C} \subset \mathcal{B}$ ,

$$\begin{aligned} \int_C E_{\mathcal{C}}[E_{\mathcal{B}}Y]dP &= \int_C E_{\mathcal{B}}YdP && \text{(definition of } E_{\mathcal{C}} \text{ as } C \in \mathcal{C}) \\ &= \int_C YdP && \text{(definition of } E_{\mathcal{B}} \text{ as } C \in \mathcal{B}). \end{aligned}$$

So  $E_{\mathcal{C}}[E_{\mathcal{B}}Y]$  satisfies the defining relation for  $E_{\mathcal{C}}Y$ . Being also  $\mathcal{C}$ -measurable, it *is*  $E_{\mathcal{C}}Y$  (a.s.). //

5'. If  $\mathcal{C} \subset \mathcal{B}$ ,  $E[E[Y|\mathcal{C}]|\mathcal{B}] = E[Y|\mathcal{C}]$  a.s.

*Proof.*  $E[Y|\mathcal{C}]$  is  $\mathcal{C}$ -measurable, so  $\mathcal{B}$ -measurable as  $\mathcal{C} \subset \mathcal{B}$ , so  $E[.|\mathcal{B}]$  has no effect on it, by 3.

*Note.* 5, 5' are the two forms of the *iterated conditional expectations property*. When conditioning on two  $\sigma$ -fields, one larger (finer), one smaller (coarser), the coarser rubs out the effect of the finer, either way round. This is also called the *coarse-averaging property*, or (Williams [W]) the *tower property*.

6. *Conditional Mean Formula.*  $E[E[Y|\mathcal{B}]] = EY$   $P$ -a.s.

*Proof.* Take  $\mathcal{C} = \{\emptyset, \Omega\}$  in 5 and use 1. //

*Example.* Check this for the bivariate normal distribution considered above.

*Note.* Compare this with the *Conditional Variance Formula* of Statistics: see e.g. SMF, IV.6, Day 9.

7. *Role of independence.* If  $Y$  is independent of  $\mathcal{B}$ ,

$$E[Y|\mathcal{B}] = EY \quad a.s.$$

*Proof.* See [W], p.88, 90, property (k).

*Note.* In the elementary definition  $P(A|B) := P(A \cap B)/P(B)$  (if  $P(B) > 0$ ), if  $A$  and  $B$  are independent (that is, if  $P(A \cap B) = P(A).P(B)$ ), then  $P(A|B) = P(A)$ : conditioning on something independent has no effect. One would expect this familiar and elementary fact to hold in this more general situation also. It does – and the proof of this rests on the proof above.

*Projections.* In Property 5 (tower property), take  $\mathcal{B} = \mathcal{C}$ :

$$E[E[X|\mathcal{C}]|\mathcal{C}] = E[X|\mathcal{C}].$$

This says that the operation of taking conditional expectation given a sub- $\sigma$ -field  $\mathcal{C}$  is *idempotent* – doing it twice is the same as doing it once. Also, taking conditional expectation is a *linear* operation (it is defined via an integral, and integration is linear). Recall from Linear Algebra that we have met such idempotent linear operations before. They are the *projections*. (Example:  $(x, y, z) \mapsto (x, y, 0)$  projects from 3-dimensional space onto the  $(x, y)$ -plane.) This view of conditional expectation as projection is useful and powerful; see e.g. [BK], [BF] or J. Neveu, *Discrete-parameter martingales* (North-Holland, 1975), I.2. It is particularly useful when one has not yet got used to conditional expectation defined measure-theoretically as above, as it gives us an alternative (and perhaps more familiar) way to think.