

M2S1 : EXERCISE SHEET 0 : SOLUTIONS

1. For events $A, B \subseteq \Omega$,

$$\omega \in A \cap B \iff \omega \in A \text{ and } \omega \in B \quad \omega \in A \cup B \iff \omega \in A \text{ or } \omega \in B \text{ or } \omega \in A \cap B$$

Hence

$$(i) \quad \omega \in E \cap \emptyset \iff \omega \in E \text{ and } \omega \in \emptyset. \text{ No such } \omega \text{ exists } \therefore E \cap \emptyset = \emptyset \\ \omega \in E \cup \emptyset \iff \omega \in E \text{ or } \omega \in \emptyset \text{ or } \omega \in E \cap \emptyset \iff \omega \in E \therefore E \cup \emptyset = E \text{ (as } (\omega \in E \cap \emptyset = \emptyset)).$$

$$(ii) \quad \omega \in E \cap \Omega \iff \omega \in E \text{ and } \omega \in \Omega \iff \omega \in E \therefore E \cap \Omega = E \\ \omega \in E \cup \Omega \iff \omega \in E \text{ or } \omega \in \Omega \text{ or } \omega \in E \cap \Omega \iff \omega \in \Omega \therefore E \cup \Omega = \Omega$$

$$(iii) \quad (E \cap F) \cup (E' \cup F') = ((E \cap F) \cup E') \cup ((E \cap F) \cup F') \\ = ((E \cup E') \cap (F \cup E')) \cup ((E \cup F') \cap (F \cup F')) \\ = (\Omega \cap (F \cup E')) \cup ((E \cup F') \cap \Omega) \\ = (F \cup E') \cup (E \cup F') \\ = \Omega$$

$$\therefore (E \cap F) \cup (E' \cup F') = \Omega \text{ so } (E \cap F)' = (E' \cup F') \text{ and } (E' \cup F')' = (E \cap F)$$

$$\text{Also, } (E \cup F)' = (E' \cap F') \text{ (by replacing } E \text{ and } F \text{ by } E' \text{ and } F')$$

$$(iv) \quad E \subseteq F \implies F = E \cup G, \text{ say, where } E \cap G = \emptyset \implies F' = (E \cup G)' = E' \cap G' \subseteq E'$$

$$(v) \quad \text{If } E \subseteq F, \text{ then } F = E \cup G, \text{ say, where } E \cap G = \emptyset \text{ so}$$

$$E \cap F = E \cap (E \cup G) = (E \cap E) \cup (E \cap G) = E \\ E \cup F = E \cup (E \cup G) = (E \cup G) = F.$$

2. An event is a subset of sample outcomes, and the number of subsets of a collection of k items is 2^k (each of the k outcomes can be either included or excluded from the subsets).

3. A collection of subsets, \mathcal{A} , of sample space Ω , say

$$\mathcal{A} = \{A_1, A_2, \dots\},$$

is a σ -algebra if

$$(I) \quad \Omega \in \mathcal{A}$$

$$(II) \quad A \in \mathcal{A} \implies A' \in \mathcal{A}$$

$$(III) \quad A_1, A_2, \dots \in \mathcal{A} \implies \bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$$

Taking these three conditions in turn we attempt to construct a minimal collection \mathcal{A}_0 of subsets that satisfies these conditions. Clearly (I) requires that $\Omega \in \mathcal{A}_0$, and (II) implies that, as $\Omega \in \mathcal{A}_0$, we require $\emptyset \in \mathcal{A}_0$. Now, if $\mathcal{A}_0 = \{\emptyset, \Omega\}$, then it can be verified that (III) holds as

$$\emptyset \cup \Omega = \Omega \in \mathcal{A}_0.$$

Hence $\mathcal{A}_0 = \{\emptyset, \Omega\}$ is the smallest σ -algebra of subsets of Ω .

4.(i) $\Omega = \{H, T\}$, so the set \mathcal{A} of all subsets of Ω , is

$$\{ \emptyset, \{H\}, \{T\}, \{H, T\} = \Omega \},$$

and it is easy to verify that \mathcal{A} is a σ -algebra by checking conditions (I), (II) and (III).

(ii) If Ω is countable, say $\Omega = \{ \omega_1, \omega_2, \dots \}$, let \mathcal{A} be the set of all subsets of Ω . An event A_i is a subset of Ω if $\omega \in A_i \implies \omega \in \Omega$, so \mathcal{A} is the set of events A_i that are countable unions of the elements of Ω .

Therefore

(I) $\Omega \in \mathcal{A}$ by construction.

(II) $A \in \mathcal{A} \implies A' \in \mathcal{A}$ (as $A \cup A' = \Omega$, and A and Ω are countable unions of elements of Ω).

(III) \mathcal{A} is closed under countable union (as the countable union of countable unions of elements of Ω is itself a countable unions of elements of Ω).

and hence \mathcal{A} is a σ -algebra.

5.(a) $\{A_k\}$ forms a partition of $\Omega = \mathbb{R}^+$ as $\omega \in \Omega \implies \omega \in A_k$ for precisely one k , and $\bigcup_{k=1}^{\infty} A_k = \Omega$.

Construct σ -algebra \mathcal{A} whose elements are countable unions of the events A_k .

A countable union of elements of the $\{A_k\}$ sequence can be described by an infinite binary sequence in which the k th term is 1 if A_k is in the countable union, and 0 otherwise. It is then straightforward to verify that the conditions (I), (II) and (III).

(I) consider the infinite binary sequence whose elements are all 1s.

(II) If $A \in \mathcal{A}$, then A' has an binary sequence representation obtained by switching 1 to 0 and 0 to 1 for each element of the A sequence. Then clearly $A' \in \mathcal{A}$.

(III) \mathcal{A} is closed under countable union, by considering the binary sequence representation.

and hence \mathcal{A} is a σ -algebra.

(b) Can construct all such intervals by taking unions and intersections of A_i events and their complements, (if we interpret $\{a_i\}$ as the interval $(a_i, a_i]$), as

$$A_i' \cap A_j = (a_i, a_j] \quad a_i < a_j$$

and

$$[a, b] = (a, b] \cup \{a\} \quad [a, b) = [a, b] \cup \{b\}' \quad (a, b) = (a, b] \cap \{b\}'$$

(ii) Straightforward to verify that conditions (I), (II), and (III) hold using the distributive and De Morgan laws