

MODULARITY LIFTING THEOREMS - NOTES FOR ARIZONA WINTER SCHOOL -  
DRAFT VERSION

TOBY GEE

CONTENTS

1. Introduction	2
1.1. Notation	3
2. Galois representations	3
2.1. Basics of Galois representations (and structure of Galois groups)	3
2.6. Local representations with $p \neq l$ : the monodromy theorem	4
2.20. Local representations with $p = l$ : $p$ -adic Hodge theory	8
2.24. Number fields	9
2.33. Sources of Galois representations	12
3. Galois deformations	13
3.1. Generalities	13
3.10. Tangent spaces	15
3.15. Deformation conditions	15
3.18. Fixing determinants	16
3.20. Global deformations with local conditions	17
3.23. Presenting global deformation rings over local lifting rings	17
3.25. Finiteness of maps between global deformation rings	20
3.27. Local deformation rings with $l = p$	21
3.29. Local deformation rings with $p \neq l$	21
3.30. Deformations of fixed type	22
3.32. Taylor–Wiles deformations	22
3.35. Taylor’s “Ihara avoidance” deformations	23

---

*Date:* April 27, 2014.

These notes are based in part on a lecture course given by Richard Taylor at Harvard University in 2009.

4. Modular and automorphic forms, and the Langlands correspondence	24
4.1. The local Langlands correspondence (and the Jacquet–Langlands correspondence)	24
4.6. Hecke operators	26
4.8. Modular forms and automorphic forms on quaternion algebras	27
4.19. Galois representations associated to automorphic representations	30
5. The Taylor–Wiles–Kisin method	32
5.2. The integral theory of automorphic forms	32
5.4. Base change	36
5.5. Patching	36
6. Level raising and lowering	45
6.1. Level raising and lowering: the Khare–Wintenberger method	45
7. The project (Arizona Winter School March 2013)	46
8. Generalisations	47
8.1. Relaxing the hypotheses	47
8.2. Further generalisations	50
References	52

## 1. INTRODUCTION

The first aim of these notes is to explain modularity/automorphy lifting theorems for two-dimensional  $p$ -adic representations, using wherever possible arguments that go over to the  $n$ -dimensional case. In particular, we use Taylor’s arguments in [Tay08] that avoid the use of Ihara’s lemma. For the most part I ignore the issues which are local at  $p$ , focusing on representations which satisfy the Fontaine–Laffaille condition.

The second aim is to explain the application of these theorems to questions of level raising and lowering for (Hilbert) modular forms, via the method of Khare–Wintenberger. This is sketched, with the details left as an exercise, to form the first part of the project.

I would like to thank Kevin Buzzard, Kestutis Cesnavicius, Jessica Fintzen, Jeff Hatley, Christian Johansson, Keenan Kidwell, Jack Lamplugh, Tom Lovering, Judith Ludwig, James Newton, Sug Woo Shin and Jack Shotton for their helpful comments on earlier versions of these notes.

**1.1. Notation.** Much of this notation will also be introduced in the text, but I have tried to collect together various definitions here, for ease of reading. Throughout these notes,  $p > 2$  is a prime greater than two. In the earlier stages of the notes, we discuss  $n$ -dimensional  $p$ -adic and mod  $p$  representations, before specialising to the case  $n = 2$ . When we do so, we assume that  $p \nmid n$ . (Of course, in the case  $n = 2$ , this follows from our assumption that  $p > 2$ .)

If  $M$  is a field, we let  $G_M$  denote its absolute Galois group. We write  $\varepsilon_p$  for the  $p$ -adic cyclotomic character. We fix an algebraic closure  $\overline{\mathbb{Q}}$  of  $\mathbb{Q}$ , and regard all algebraic extensions of  $\mathbb{Q}$  as subfields of  $\overline{\mathbb{Q}}$ . For each prime  $p$  we fix an algebraic closure  $\overline{\mathbb{Q}}_p$  of  $\mathbb{Q}_p$ , and we fix an embedding  $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$ . In this way, if  $v$  is a finite place of a number field  $F$ , we have a homomorphism  $G_{F_v} \hookrightarrow G_F$ . We also fix an embedding  $\overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$ . If  $L$  is a local field, we denote its residue field by  $k(L)$ .

We normalise the definition of Hodge–Tate weights so that all the Hodge–Tate weights of the  $p$ -adic cyclotomic character  $\varepsilon_p$  are  $-1$ .

We let  $\zeta_p$  be a primitive  $p$ th root of unity.

## 2. GALOIS REPRESENTATIONS

**2.1. Basics of Galois representations (and structure of Galois groups).** Let  $K'/K$  be a (not necessarily finite) normal and separable extension of fields. Then the Galois group  $\text{Gal}(K'/K)$  is the group

$$\{\sigma \in \text{Aut}(K') : \sigma|_K = \text{id}_K\}.$$

This has a natural topology, making it a compact Hausdorff totally disconnected topological group; equivalently, it is a profinite group. This can be expressed by the topological isomorphism

$$\text{Gal}(K'/K) \cong \varprojlim_{K''/K \text{ finite normal}} \text{Gal}(K''/K),$$

where the finite groups  $\text{Gal}(K''/K)$  have the discrete topology.

Then Galois theory gives a bijective correspondence between intermediate fields  $K' \supset K'' \supset K$  and closed subgroups  $H \subset \text{Gal}(K'/K)$ , with  $K''$  corresponding to  $\text{Gal}(K'/K'')$  and  $H$  corresponding to  $K^H$ .

Fix a separable closure  $\overline{K}$  of  $K$ , and write  $G_K := \text{Gal}(\overline{K}/K)$ . Let  $L$  be a topological field; then a *Galois representation* is a continuous homomorphism  $\rho : G_K \rightarrow \text{GL}_n(L)$  for some  $n$ . The nature of these representations depends on the topology on  $L$ . For example, if  $L$  has the discrete topology, then the image of  $\rho$  is finite, and  $\rho$  factors through a finite Galois group  $\text{Gal}(K''/K)$ .

**2.2. Exercise.** If  $L = \mathbb{C}$  with the usual topology, then  $\rho(G_K)$  is finite, and  $\rho$  is conjugate to a representation valued in  $\text{GL}_n(\overline{\mathbb{Q}})$ .

On the other hand, if  $L/\mathbb{Q}_p$  is a finite extension with the  $p$ -adic topology, then there can be examples with infinite image. The rest of this course will be concerned with these  $p$ -adic representations. For example, if  $p \neq \text{char } K$ , we have the  $p$ -adic cyclotomic character  $\varepsilon_p : G_K \rightarrow \mathbb{Z}_p^\times$ , which is uniquely determined by the requirement that if  $\sigma \in G_K$  and  $\zeta \in \bar{K}$  with  $\zeta^{p^n} = 1$  for some  $n$ , then  $\sigma(\zeta) = \zeta^{\varepsilon_p(\sigma) \pmod{p^n}}$ .

2.3. *Fact.* If  $L/\mathbb{Q}_p$  is an algebraic extension, and  $\rho : G_K \rightarrow \text{GL}_n(L)$  is a continuous representation, then  $\rho(G_K) \subseteq \text{GL}_n(M)$  for some  $L \supset M \supset \mathbb{Q}_p$  with  $M/\mathbb{Q}_p$  finite.

*Proof.* This follows from the Baire category theorem; see e.g. the proof of Corollary 5 of [Dic01] for the details.  $\square$

2.4. *Exercise.* If  $L/\mathbb{Q}_p$  is an algebraic extension, and  $\rho : G_K \rightarrow \text{GL}_n(L)$  is a continuous representation, then  $\rho$  is conjugate to a representation in  $\text{GL}_n(\mathcal{O}_L)$ .

Any finite-dimensional Galois representation has a Jordan–Hölder sequence, and thus a well-defined semisimplification.

2.5. *Fact.* Two Galois representations  $\rho, \rho' : G_K \rightarrow \text{GL}_n(L)$  have isomorphic semisimplifications if and only if  $\rho(g), \rho'(g)$  have the same characteristic polynomials for each  $g \in G_K$ . If  $\text{char } L = 0$  (or indeed if  $\text{char } L > n$ ), then this is equivalent to  $\text{tr } \rho(g) = \text{tr } \rho'(g)$  for all  $g \in G_K$ .

*Proof.* This is the Brauer–Nesbitt theorem, cf. [CR62, 30.16]  $\square$

As a corollary of the previous exercise and fact, we see that  $p$ -adic representations have well-defined semi-simplified reductions modulo  $p$ . Indeed, given  $\rho : G_K \rightarrow \text{GL}_n(L)$  with  $L/\mathbb{Q}_p$  algebraic, we may conjugate  $\rho$  to be valued in  $\text{GL}_n(\mathcal{O}_L)$ , reduce modulo the maximal ideal and semisimplify to get a semisimple representation  $\bar{\rho} : G_K \rightarrow \text{GL}_n(k(L))$ , whose characteristic polynomials are determined by those of  $\rho$ .

[We really do have to semisimplify here; to see why, think about the reductions modulo  $p$  of the matrices  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$ .]

2.6. **Local representations with  $p \neq l$ : the monodromy theorem.** In this section we will let  $K/\mathbb{Q}_l$  be a finite extension, for some prime  $l \neq p$ . In order to study the representations of  $G_K$ , we firstly recall something of the structure of  $G_K$  itself; cf. [Ser79] for further details. Let  $\varpi$  be a uniformiser of  $\mathcal{O}_K$ , and let  $\text{val}_K : K^\times \rightarrow \mathbb{Z}$  be the  $\varpi$ -adic valuation. Let  $|\cdot|_K := (\#k)^{-\text{val}_K(\cdot)}$  be the corresponding norm. The action of  $G_K$  on  $K$  preserves  $\text{val}_K$ , and thus induces an action on  $k$ , so that we have a homomorphism  $G_K \rightarrow G_k$ , and in fact a short exact sequence

$$0 \rightarrow I_K \rightarrow G_K \rightarrow G_k \rightarrow 0$$

defining the inertia subgroup  $I_K$ . We let  $\text{Frob}_K = \text{Frob}_k \in G_k$  be the geometric Frobenius element, a generator of  $G_k \cong \widehat{\mathbb{Z}}$ .

Then we define the Weil group  $W_K$  via the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & I_K & \longrightarrow & G_K & \longrightarrow & G_k \longrightarrow 0 \\ & & \parallel & & \uparrow & & \uparrow \\ 0 & \longrightarrow & I_K & \longrightarrow & W_K & \longrightarrow & \text{Frob}_k^{\mathbb{Z}} \longrightarrow 0 \end{array}$$

so that  $W_K$  is the subgroup of  $G_K$  consisting of elements which map to an integral power of the Frobenius in  $G_k$ . The group  $W_K$  is a topological group, but its topology is not the subspace topology of  $G_K$ ; rather, the topology is determined by decreeing that  $I_K$  is open, and has its usual topology.

Let  $K^{\text{ur}} = \overline{K}^{I_K}$  be the maximal unramified extension of  $K$ , and let  $K^{\text{tame}} = \cup_{(n,l)=1} K^{\text{ur}}(\omega_K^{1/n})$  be the maximal tamely ramified extension. Then the wild inertia subgroup  $P_K := \text{Gal}(\overline{K}/K^{\text{tame}})$  is the unique Sylow pro- $l$  subgroup of  $I_K$ . Let  $\zeta = (\zeta_n)_{(n,l)=1}$  be a compatible system of primitive roots of unity (i.e.  $\zeta_{ab}^a = \zeta_b$ ). Then we have a character

$$t_\zeta : I_K/P_K \xrightarrow{\sim} \prod_{p \neq l} \mathbb{Z}_p,$$

defined by

$$\frac{\sigma(\omega_K^{1/n})}{\omega_K^{1/n}} = \zeta_n^{(t_\zeta(\sigma) \pmod{n})}.$$

**2.7. Exercise.** Any other compatible system of roots of unity is of the form  $\zeta^u$  for some  $u \in \prod_{p \neq l} \mathbb{Z}_p^\times$ , and we have  $t_{\zeta^u} = u^{-1}t_\zeta$ .

If  $\sigma \in W_K$ , then  $t_\zeta(\sigma\tau\sigma^{-1}) = \varepsilon(\sigma)t_\zeta(\tau)$ , where  $\varepsilon$  is the cyclotomic character. We let  $t_{\zeta,p}$  be the composite of  $t_\zeta$  and the projection to  $\mathbb{Z}_p$ .

Local class field theory is summarised in the following statement.

**2.8. Theorem.** Let  $W_K^{\text{ab}}$  denote the group  $W_K/\overline{[W_K, W_K]}$ . Then there are unique isomorphisms  $\text{Art}_K : K^\times \xrightarrow{\sim} W_K^{\text{ab}}$  such that

- (1) if  $K'/K$  is a finite extension, then  $\text{Art}_{K'} = \text{Art}_K \circ N_{K'/K}$ , and
- (2) we have a commutative square

$$\begin{array}{ccc} K^\times & \xrightarrow{\text{Art}_K} & W_K^{\text{ab}} \\ \downarrow \text{val}_K & & \downarrow \\ \mathbb{Z} & \longrightarrow & \text{Frob}_K^{\mathbb{Z}} \end{array}$$

where the bottom arrow is the isomorphism sending  $a \mapsto \text{Frob}_K^a$ .

The irreducible representations of the group  $W_K^{\text{ab}}$  are just the characters of  $W_K$ , and local class field theory gives a simple description of them, as representations of  $K^\times = \text{GL}_1(K)$ . The

local Langlands correspondence for  $\mathrm{GL}_n$  (see Section 4.1) is a kind of  $n$ -dimensional generalisation of this, giving a description of the  $n$ -dimensional representations of  $W_K$  in terms of the representation theory of  $\mathrm{GL}_n(K)$ .

**2.9. Definition.** Let  $L$  be a field of characteristic 0. A *representation* of  $W_K$  over  $L$  is a representation (on a finite-dimensional  $L$ -vector space) which is continuous if  $L$  has the discrete topology (i.e. a representation with open kernel).

A *Weil–Deligne* representation of  $W_K$  on a finite-dimensional  $L$ -vector space  $V$  is a pair  $(r, N)$  consisting of a representation  $r : W_K \rightarrow \mathrm{GL}(V)$ , and an endomorphism  $N \in \mathrm{End}(V)$  such that for all  $\sigma \in W_K$ ,

$$r(\sigma)Nr(\sigma)^{-1} = (\#\kappa)^{-v_K(\sigma)}N,$$

where  $v_K : W_K \rightarrow \mathbb{Z}$  is determined by  $\sigma|_{K^{\mathrm{ur}}} = \mathrm{Frob}_K^{v_K(\sigma)}$ .

**2.10. Remark.** (1) Since  $I_K$  is compact and open in  $W_K$ , if  $r$  is a representation of  $W_K$  then  $r(I_K)$  is finite.

(2)  $N$  is necessarily nilpotent.

**2.11. Exercise.** (1) Show that if  $(r, V)$  is a representation of  $W_K$  and  $m \geq 1$  then the following defines a Weil–Deligne representation  $\mathrm{Sp}_m(r)$  with underlying vector space  $V^m$ : we let  $W_K$  act via

$$r| \mathrm{Art}_K^{-1}|_K^{m-1} \oplus r| \mathrm{Art}_K^{-1}|_K^{m-2} \oplus \cdots \oplus r,$$

and let  $N$  induce an isomorphism from  $r| \mathrm{Art}_K^{-1}|_K^{i-1}$  to  $r| \mathrm{Art}_K^{-1}|_K^i$  for each  $i < m - 1$ , and be 0 on  $r| \mathrm{Art}_K^{-1}|_K^{m-1}$ .

(2) Show that every Weil–Deligne representation  $(r, V)$  for which  $r$  is semisimple is isomorphic to a direct sum of representations  $\mathrm{Sp}_{m_i}(r_i)$ .

(3) Show that if  $(r, V, N)$  is a Weil–Deligne representation of  $W_K$ , and  $K'/K$  is a finite extension, then  $(r|_{W_{K'}}, V, N)$  is a Weil–Deligne representation of  $W_{K'}$ .

(4) Suppose that  $r$  is a representation of  $W_K$ . Show that if  $\sigma \in W_K$  then for some positive integer  $n$ ,  $r(\sigma^n)$  is in the centre of  $r(W_K)$ .

(5) Assume further that  $\sigma \notin I_K$ . Show that for any  $\tau \in W_K$  there exists  $n \in \mathbb{Z}$  and  $m > 0$  such that  $r(\sigma^n) = r(\tau^m)$ .

(6) Show that for a representation  $r$  of  $W_K$ , the following conditions are equivalent:

- (a)  $r$  is semisimple.
- (b)  $r(\sigma)$  is semisimple for all  $\sigma \in W_K$ .
- (c)  $r(\sigma)$  is semisimple for some  $\sigma \notin I_K$ .

(7) Let  $(r, N)$  be a Weil–Deligne representation of  $W_K$ . Set  $\tilde{r}(\sigma) = r(\sigma)^{\mathrm{ss}}$ , the semisimplification of  $r(\sigma)$ . Prove that  $(\tilde{r}, N)$  is also a Weil–Deligne representation of  $W_K$ .

**2.12. Definition.** We say that a Weil–Deligne representation  $(r, N)$  is *Frobenius-semisimple* if  $r$  is semisimple. With notation as above, we say that  $(\tilde{r}, N)$  is the *Frobenius semisimplification* of  $(r, N)$ .

2.13. **Definition.** If  $L$  is an algebraic extension of  $\mathbb{Q}_p$ , then we say that an element  $A \in \mathrm{GL}_n(L)$  is *bounded* if it has determinant in  $\mathcal{O}_L^\times$ , and characteristic polynomial in  $\mathcal{O}_L[X]$ .

2.14. *Exercise.*  $A$  is bounded if and only if it stabilises an  $\mathcal{O}_L$ -lattice in  $L^n$ .

2.15. **Definition.** Let  $L$  be an algebraic extension of  $\mathbb{Q}_p$ . Then we say that  $r$  is *bounded* if  $r(\sigma)$  is bounded for all  $\sigma \in W_K$ .

2.16. *Exercise.* Show  $r$  is bounded if and only if  $r(\sigma)$  is bounded for some  $\sigma \notin I_K$ .

The reason for all of these definitions is the following theorem, which in practice gives us a rather concrete classification of the  $p$ -adic representations of  $G_K$ .

2.17. **Proposition.** (*Grothendieck's monodromy theorem*) Suppose that  $l \neq p$ , that  $K/\mathbb{Q}_l$  is finite, and that  $V$  is a finite-dimensional  $L$ -vector space, with  $L$  an algebraic extension of  $\mathbb{Q}_p$ . Fix  $\varphi \in W_K$  a lift of  $\mathrm{Frob}_K$  and a compatible system  $(\zeta_n)_{(n,l)=1}$  of primitive roots of unity. If  $\rho : G_K \rightarrow \mathrm{GL}(V)$  is a continuous representation then there is a finite extension  $K'/K$  and a uniquely determined nilpotent  $N \in \mathrm{End}(V)$  such that for all  $\sigma \in I_{K'}$ ,

$$\rho(\sigma) = \exp(Nt_{\zeta,p}(\sigma)).$$

For all  $\sigma \in W_K$ , we have  $\rho(\sigma)N\rho(\sigma)^{-1} = \#\kappa^{-v(\sigma)}N$ . In fact, we have an equivalence of categories  $\mathrm{WD} = \mathrm{WD}_{\zeta,\varphi}$  from the category of continuous representations of  $G_K$  on finite-dimensional  $L$ -vector spaces to the category of bounded Weil–Deligne representations on finite-dimensional  $L$ -vector spaces, taking

$$\rho \mapsto (V, r, N), \quad r(\tau) := \rho(\tau) \exp(-t_{\zeta,p}(\varphi^{-v_K(\tau)}\tau)N).$$

The functors  $\mathrm{WD}_{\zeta',\varphi'}$  and  $\mathrm{WD}_{\zeta,\varphi}$  are naturally isomorphic.

2.18. *Remark.* Note that since  $N$  is nilpotent, the exponential here is just a polynomial - there are no convergence issues!

The proof is contained in the following exercise.

- 2.19. *Exercise.*
- (1) By Exercise 2.4 there is a  $G_K$ -stable  $\mathcal{O}_L$ -lattice  $\Lambda \subset V$ . Show that if  $G_{K'}$  is the kernel of the induced map  $G_K \rightarrow \mathrm{Aut}(\Lambda/p\Lambda)$ , then  $K'/K$  is a finite extension, and  $\rho(G_{K'})$  is pro- $p$ . Show that  $\rho|_{I_{K'}}$  factors through  $t_{\zeta,p} : I_{K'} \rightarrow \mathbb{Z}_p$ .
  - (2) Choose  $\sigma \in I_{K'}$  such that  $t_{\zeta,p}(\sigma)$  topologically generates  $t_{\zeta,p}(I_{K'})$ . By considering the action of conjugation by  $\varphi$ , show that the eigenvalues of  $\rho(\sigma)$  are all  $p$ -power roots of unity. Hence show that one may make a further finite extension  $K''/K'$  such that the elements of  $\rho(I_{K''})$  are all unipotent.
  - (3) Deduce the existence of a unique nilpotent  $N \in \mathrm{End}(V)$  such that for all  $\sigma \in I_{K''}$ ,  $\rho(\sigma) = \exp(Nt_{\zeta,p}(\sigma))$ . [Hint: use the logarithm map (why are there no convergence issues?).]
  - (4) Complete the proof of the proposition, by showing that  $(r, N)$  is a Weil–Deligne representation. Where does the condition that  $r$  is bounded come in?

One significant advantage of Weil–Deligne representations over Galois representations is that there are no subtle topological issues: the topology on the Weil–Deligne representation is the discrete topology. This allows one to describe representations in a way that is “independent of  $L$ ”, and is necessary to make sense of the notion of a compatible system of Galois representations (or at least to make sense of it at places at which the Galois representation is ramified).

**2.20. Local representations with  $p = l$ :  $p$ -adic Hodge theory.** The case  $l = p$  is far more complicated than the case  $l \neq p$ , largely because wild inertia can act in a highly nontrivial fashion, so there is no simple analogue of Grothendieck’s monodromy theorem. (There is still an analogue, though, it’s just much harder to state and prove, and doesn’t apply to all  $p$ -adic Galois representations.) The study of representations  $G_K \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_p)$  with  $K/\mathbb{Q}_p$  finite is called  *$p$ -adic Hodge theory*, a subject largely developed by Fontaine in the 1980s. An excellent introduction to the subject can be found in [BC], and the standard reference is [Fon94]. We will content ourselves with some terminology, some definitions, and some remarks intended to give intuition and motivation.

Fix  $K/\mathbb{Q}_p$  finite. In some sense, “most”  $p$ -adic Galois representations  $G_K \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_p)$  will not be relevant for us, because they do not arise in geometry, or in the Galois representations associated to automorphic representations. Instead, there is a hierarchy of classes of representations

$$\{\text{crystalline}\} \subsetneq \{\text{semistable}\} \subsetneq \{\text{de Rham}\} \subsetneq \{\text{Hodge–Tate}\}.$$

For any of these classes  $X$ , we say that  $\rho$  is *potentially*  $X$  if there is a finite extension  $K'/K$  such that  $\rho|_{G_{K'}}$  is  $X$ . A representation is potentially de Rham if and only if it is de Rham, and potentially Hodge–Tate if and only if it is Hodge–Tate; the corresponding statements for crystalline and semistable representations are false, as we will see concretely in the case  $n = 1$  later on. The  $p$ -adic analogue of Grothendieck’s monodromy theorem is the following deep theorem of Berger.

**2.21. Theorem.** *(The  $p$ -adic monodromy theorem) A representation is de Rham if and only if it is potentially semistable.*

The notion of a de Rham representation is designed to capture the representations arising in geometry; it does so by the following result of Tsuji (building on the work of many people).

**2.22. Theorem.** *If  $X/K$  is a smooth projective variety, then each  $H_{\text{ét}}^i(X \times_K \overline{K}, \overline{\mathbb{Q}}_p)$  is a de Rham representation.*

Similarly, the definitions of crystalline and semistable are designed to capture the notions of good and semistable reduction, and one has

**2.23. Theorem.** *If  $X/K$  is a smooth projective variety with good (respectively, semistable) reduction, then each  $H_{\text{ét}}^i(X \times_K \overline{K}, \overline{\mathbb{Q}}_p)$  is a crystalline (respectively, semistable) representation.*



Thus the  $p$ -adic monodromy theorem can be thought of as a Galois-theoretic incarnation of Grothendieck's semistable reduction theorem.

The case that  $n = 1$  is particularly simple, as we now explain. In this case, every semistable character is crystalline, and the de Rham characters are exactly the Hodge–Tate characters. In the case  $K = \mathbb{Q}_p$ , these are precisely the characters whose restrictions to inertia are of the form  $\psi \varepsilon_p^n$  where  $\psi$  has finite order and  $n \in \mathbb{Z}$ , while the crystalline characters are those for which  $\psi$  is unramified. A similar description exists for general  $K$ , with  $\varepsilon_p^n$  replaced by a product of so-called *Lubin–Tate characters*. In fact (c.f. Exercise 6.4.3 of [BC]), a character  $\chi : G_K \rightarrow \overline{\mathbb{Q}_p}^\times$  is de Rham if and only if there is an open subgroup  $U$  of  $K^\times$  and an integer  $n_\tau$  for each  $\tau : K \hookrightarrow \overline{\mathbb{Q}_p}^\times$  such that  $(\chi \circ \text{Art}_K)(\alpha) = \prod_\tau \tau(\alpha)^{-n_\tau}$  for each  $\alpha \in U$ , and it is crystalline if and only if we can take  $U = \mathcal{O}_K^\times$ .

As soon as  $n > 1$ , there are non-crystalline semistable representations, and non-de Rham Hodge–Tate representations. A useful heuristic when comparing to the  $l \neq p$  case is that crystalline representations correspond to unramified representations, semistable representations correspond to representations for which inertia acts unipotently, and de Rham representations correspond to all representations.

Suppose that  $\rho : G_K \rightarrow \text{GL}_n(\overline{\mathbb{Q}_p})$  is a Hodge–Tate representation. Then for each  $\tau : K \hookrightarrow \overline{\mathbb{Q}_p}$  there is a multiset of  $\tau$ -labeled *Hodge–Tate weights*  $\text{HT}_\tau(\rho)$  associated to  $\rho$ ; this is a multiset of integers, and in the case of a de Rham character  $\chi$  as above,  $\text{HT}_\tau(\chi) = n_\tau$ . In particular, the  $p$ -adic cyclotomic character  $\varepsilon_p$  has all Hodge–Tate weights equal to  $-1$ . If  $K'/K$  is a finite extension, and  $\tau' : K' \hookrightarrow \overline{\mathbb{Q}_p}$  extends  $\tau : K \hookrightarrow \overline{\mathbb{Q}_p}$ , then  $\text{HT}_{\tau'}(\rho|_{G_{K'}}) = \text{HT}_\tau(\rho)$ .

If furthermore  $\rho$  is potentially semistable (equivalently, de Rham) then a construction of Fontaine associates a Weil–Deligne representation  $\text{WD}(\rho) = (r, N)$  of  $W_K$  to  $\rho$ . If  $K'/K$  is a finite extension, then  $\text{WD}(\rho|_{G_{K'}}) = (r|_{W_{K'}}, N)$ . It is known that  $\rho$  is semistable if and only if  $r$  is unramified, and that  $\rho$  is crystalline if and only if  $r$  is unramified and  $N = 0$ . Thus  $\rho$  is potentially crystalline if and only if  $N = 0$ .

**2.24. Number fields.** We now consider the case that  $K$  is a number field (that is, a finite extension of  $\mathbb{Q}$ ). If  $v$  is a finite place of  $K$ , we let  $K_v$  denote the completion of  $K$  at  $v$ . If  $K'/K$  is a finite Galois extension, then  $\text{Gal}(K'/K)$  transitively permutes the places of  $K'$  above  $v$ ; if we choose one such place  $w$ , then we define the *decomposition group*

$$\text{Gal}(K'/K)_w := \{\sigma \in \text{Gal}(K'/K) \mid w\sigma = w\}.$$

Then we have a natural isomorphism  $\text{Gal}(K'/K)_w \xrightarrow{\sim} \text{Gal}(K'_w/K_v)$ , and since  $\text{Gal}(K'/K)_{w\sigma} = \sigma^{-1}\text{Gal}(K'/K)_w\sigma$ , we see that the definition extends to general algebraic extensions, and in particular we have an embedding  $G_{K_v} \hookrightarrow G_K$  which is well-defined up to conjugacy (alternatively, up to a choice of embedding  $\overline{K} \hookrightarrow \overline{K}_v$ ). (Note that you need to be slightly careful with

taking completions in the case that  $K'/K$  is infinite, as then the extension  $K'_w/K_v$  need not be algebraic, we can for example define  $\text{Gal}(K'_w/K_v)$  to be the group of continuous automorphisms of  $K'_w$  which fix  $K_v$  pointwise.)

If  $K'/K$  is Galois and unramified at  $v$ , and  $w$  is a place of  $K'$  lying over  $v$ , then we define

$$\text{Frob}_w := \text{Frob}_{K_v} \in \text{Gal}(K'_w/K_v) \xrightarrow{\sim} \text{Gal}(K'/K)_w \hookrightarrow \text{Gal}(K'/K).$$

We have  $\text{Frob}_{w\sigma} = \sigma^{-1} \text{Frob}_w \sigma$ , and thus a well-defined conjugacy class  $[\text{Frob}_v] = \{\text{Frob}_w\}_{w|v}$  in  $\text{Gal}(K'/K)$ .

**2.25. Fact.** (Chebotarev density theorem) If  $K'/K$  is a Galois extension which is unramified outside of a finite set  $S$  of places of  $K$ , then the union of the conjugacy classes  $[\text{Frob}_v]$ ,  $v \notin S$  is dense in  $\text{Gal}(K'/K)$ .

We briefly recall the statement of global class field theory. Let  $\mathbb{A}_K$  denote the adèles of  $K$ , and write  $K_\infty = \prod_{v|\infty} K_v$ . Let  $K^{\text{ab}} = \overline{K}^{[G_K, G_K]}$  be the maximal abelian extension of  $K$ . Then there is a homomorphism  $\text{Art}_K : \mathbb{A}_K^\times / (K_\infty^\times)^\circ \rightarrow \text{Gal}(K^{\text{ab}}/K)$ , defined in the following way: for each finite place  $v$  of  $K$ , the restriction of  $\text{Art}_K$  to  $K_v^\times$  agrees with the local Artin maps  $\text{Art}_{K_v}$ , and similarly at the infinite places, it agrees with the obvious isomorphisms  $\text{Art}_{K_v} : K_v^\times / (K_v^\times)^\circ \xrightarrow{\sim} \text{Gal}(\overline{K}_v/K_v)$ . Then global class field theory states that  $\text{Art}_K$  induces an isomorphism

$$\text{Art}_K : \mathbb{A}_K^\times / \overline{K^\times (K_\infty^\times)^\circ} \xrightarrow{\sim} \text{Gal}(K^{\text{ab}}/K).$$

The global Galois representations that we will care about are those that Fontaine and Mazur call *geometric*. Let  $L/\mathbb{Q}_p$  be an algebraic extension.

**2.26. Definition.** A continuous representation  $\rho : G_K \rightarrow \text{GL}_n(L)$  is *geometric* if it is unramified outside of a finite set of places of  $K$ , and if for each place  $v|p$ ,  $\rho|_{G_{K_v}}$  is de Rham.

**2.27. Remark.** It is known that both conditions are necessary; that is, there are examples of representations which are unramified outside of a finite set of places of  $K$  but not de Rham at places lying over  $p$ , and examples of representations which are de Rham at all places lying over  $p$ , but are ramified at infinitely many primes.

In practice (and conjecturally always), geometric Galois representations arise as part of a *compatible system* of Galois representations. There are a number of different definitions of a compatible system in the literature, all of which are conjecturally equivalent (although proving the equivalence of the definitions is probably very hard). The following definition, taken from [BLGGT10], seems to incorporate the minimal assumptions under which one can hope to employ automorphy lifting theorems to study a compatible system.

**2.28. Definition.** Suppose that  $K$  and  $M$  are number fields, that  $S$  is a finite set of primes of  $K$  and that  $n$  is a positive integer. By a *weakly compatible system* of  $n$ -dimensional  $p$ -adic representations of  $G_K$  defined over  $M$  and unramified outside  $S$  we mean a family of continuous

semisimple representations

$$r_\lambda : G_K \longrightarrow \mathrm{GL}_n(\overline{M}_\lambda),$$

where  $\lambda$  runs over the finite places of  $M$ , with the following properties.

- If  $v \notin S$  is a finite place of  $K$ , then for all  $\lambda$  not dividing the residue characteristic of  $v$ , the representation  $r_\lambda$  is unramified at  $v$  and the characteristic polynomial of  $r_\lambda(\mathrm{Frob}_v)$  lies in  $M[X]$  and is independent of  $\lambda$ .
- Each representation  $r_\lambda$  is de Rham at all places above the residue characteristic of  $\lambda$ , and in fact crystalline at any place  $v \notin S$  which divides the residue characteristic of  $\lambda$ .
- For each embedding  $\tau : K \hookrightarrow \overline{M}$  the  $\tau$ -Hodge–Tate numbers of  $r_\lambda$  are independent of  $\lambda$ .

2.29. *Remark.* By the Chebotarev density theorem, each  $r_\lambda$  is determined by the characteristic polynomials of the  $r_\lambda(\mathrm{Frob}_v)$  for  $v \notin S$ , and in particular the compatible system is determined by a single  $r_\lambda$ . Note that for a general element  $\sigma \in G_K$ , there will be no relationship between the characteristic polynomials of the  $r_\lambda(\sigma)$  as  $\lambda$  varies (and they won't even lie in  $M[X]$ , so there will be no way of comparing them).

There are various other properties one could demand; for example, we have the following definition (again following [BLGGT10], although we have slightly strengthened the definition made there by allowing  $\lambda$  to divide the residue characteristic of  $v$ ).

2.30. **Definition.** We say that a weakly compatible system is *strictly compatible* if for each finite place  $v$  of  $K$  there is a Weil–Deligne representation  $\mathrm{WD}_v$  of  $W_{K_v}$  over  $\overline{M}$  such that for each place  $\lambda$  of  $M$  and every  $M$ -linear embedding  $\zeta : \overline{M} \hookrightarrow \overline{M}_\lambda$  we have  $\zeta \mathrm{WD}_v \cong \mathrm{WD}(r_\lambda|_{G_{K_v}})^{\mathrm{F-ss}}$ .

Conjecturally, every weakly compatible system is strictly compatible, and even satisfies further properties, such as purity (c.f. Section 5 of [BLGGT10]). We also have the following consequence of the Fontaine–Mazur conjecture and standard conjectures on the étale cohomology of algebraic varieties over number fields.

2.31. **Conjecture.** *Any semisimple geometric representation  $G_K \rightarrow \mathrm{GL}_n(L)$  is part of a strictly compatible system of Galois representations.*

In practice, most progress on understanding these conjectures has been made by using automorphy lifting theorems to prove special cases of the following conjecture.

2.32. **Conjecture.** *Any weakly compatible system of Galois representations is strictly compatible, and is in addition automorphic, in the sense that there is an algebraic automorphic representation (in the sense of [Clo90])  $\pi$  of  $\mathrm{GL}_n(\mathbb{A}_K)$  with the property that  $\mathrm{WD}_v(\rho) \cong \mathrm{rec}(\pi_v) |\det|^{(1-n)/2}$  for each finite place  $v$  of  $K$ , where  $\mathrm{rec}$  is the local Langlands correspondence as in Section 4.1 below.*

**2.33. Sources of Galois representations.** The main source (and conjecturally the only source) of compatible systems of Galois representations is the étale cohomology of algebraic varieties. We have the following result, whose proof is well beyond the scope of this course.

**2.34. Theorem.** *Let  $K$  be a number field, and let  $X/K$  be a smooth projective variety. Then for any  $i, j$ , the  $H_{\text{ét}}^i(X \times_K \bar{K}, \mathbf{Q}_p)^{\text{ss}}(j)$  (the  $(j)$  denoting a Tate twist) form a weakly compatible system.*

**2.35. Remark.** Conjecturally, it is a strictly compatible system, and there is no need to semisimplify the representations. Both of these properties are known if  $X$  is an abelian variety.

**2.36. Conjecture.** *(The Fontaine–Mazur conjecture, [FM95]) Any irreducible geometric representation  $\rho : G_K \rightarrow \text{GL}_n(\bar{\mathbf{Q}}_p)$  is (the extension of scalars to  $\bar{\mathbf{Q}}_p$  of) a subquotient of a representation arising from étale cohomology as in Theorem 2.34.*

**2.37. Remark.** Conjecture 2.36, together with the expectation expressed in Remark 2.35, implies Conjecture 2.31. The *Fontaine–Mazur–Langlands conjecture* is a somewhat ill-defined conjecture, which is essentially the union of Conjectures 2.31 and 2.32, expressing the expectation that an irreducible geometric Galois representation is automorphic.

When  $n = 1$ , all of these conjectures are essentially known, as we will now explain. For  $n > 1$ , we know very little (although the situation when  $K = \mathbf{Q}$  and  $n = 2$  is pretty good), and the main results that are known are as a consequence of automorphy lifting theorems (as discussed in this course) and of potential automorphy theorems (which are not discussed in this course, but should be accessible given the material we develop here; for a nice introduction, see [Buz12]).

**2.38. Definition.** A *grossencharacter* is a continuous character  $\chi : \mathbb{A}_K^\times / K^\times \rightarrow \mathbf{C}^\times$ . We say that  $\chi$  is *algebraic* (or “type  $A_0$ ”) if for each  $\tau : K \hookrightarrow \mathbf{C}$  there is an integer  $n_\tau$ , such that for each  $\alpha \in (K_\infty^\times)^\circ$ , we have  $\chi(\alpha) = \prod_\tau (\tau\alpha)^{-n_\tau}$ .

**2.39. Definition.** Let  $L$  be a field of characteristic zero such that for each embedding  $\tau : K \hookrightarrow \bar{L}$ , we have  $\tau(K) \subseteq L$ . Then an *algebraic character*  $\chi_0 : \mathbb{A}_K^\times \rightarrow \bar{L}^\times$  is a character with open kernel such that for each  $\tau : K \hookrightarrow L$  there is an integer  $n_\tau$  with the property that for all  $\alpha \in K^\times$ , we have  $\chi_0(\alpha) = \prod_\tau (\tau\alpha)^{n_\tau}$ .

**2.40. Exercise.** Show that if  $\chi_0$  is an algebraic character, then  $\chi_0$  takes values in some number field. [Hint: show that  $\mathbb{A}_K^\times / (K^\times \ker \chi_0)$  is finite, and that  $\chi_0(K^\times \ker \chi_0)$  is contained in a number field.]

**2.41. Theorem.** *Let  $E$  be a number field containing the normal closure of  $K$ . Fix embeddings  $\iota_\infty : \bar{E} \hookrightarrow \mathbf{C}$ ,  $\iota_p : \bar{E} \hookrightarrow \bar{\mathbf{Q}}_p$ . Then the following are in natural bijection.*

- (1) Algebraic characters  $\chi_0 : \mathbb{A}_K^\times \rightarrow \bar{E}^\times$ .
- (2) Algebraic grossencharacters  $\chi : \mathbb{A}_K^\times / K^\times \rightarrow \mathbf{C}^\times$ .
- (3) Continuous representations  $\rho : G_K \rightarrow \bar{\mathbf{Q}}_p^\times$  which are de Rham at all  $v|p$ .

(4) *Geometric representations*  $\rho : G_K \rightarrow \overline{\mathbb{Q}}_p^\times$ .

2.42. *Exercise.* Prove Theorem 2.41 as follows (see e.g. Section 1 of [Far] for more details). Use the non-trivial fact that if  $v|p$  is a place of  $K$ , then a representation  $\rho_v : G_{K_v} \rightarrow \overline{\mathbb{Q}}_p^\times$  is de Rham if and only if there is an open subgroup  $U \subseteq K_v^\times$  and an integer  $n_\tau$  for each  $\tau : K_v \hookrightarrow \overline{\mathbb{Q}}_p$  such that  $(\rho_v \circ \text{Art}_{K_v})(\alpha) = \prod_\tau \tau(\alpha)^{n_\tau}$  for each  $\alpha \in U$ . Use this, together with global class field theory, to show that (3)  $\implies$  (4).

For the correspondence between (1) and (2), show that we can pair up  $\chi_0$  and  $\chi$  by

$$\chi(\alpha) = \iota_\infty \left( \chi_0(\alpha) \prod_{\tau: K \hookrightarrow \mathbb{C}} \tau(\alpha_\infty)^{-n_{\iota_\infty^{-1}\tau}} \right).$$

For the correspondence between (1) and (3), show that we can pair up  $\chi_0$  and  $\rho$  by

$$(\rho \circ \text{Art}_K)(\alpha) = \iota_p \left( \chi_0(\alpha) \prod_{\tau: K \hookrightarrow \overline{\mathbb{Q}}_p} \tau(\alpha_p)^{-n_{\iota_p^{-1}\tau}} \right).$$

### 3. GALOIS DEFORMATIONS

There are a number of good introductions to the material in this section, and for the most part we will simply give basic definitions and motivation, and refer elsewhere for proofs. In particular, [Maz97] is a very nice introduction to Galois deformations (although slightly out of date, as it does not treat liftings/framed deformations), and [Boe] is a thorough modern treatment.

**3.1. Generalities.** Take  $L/\mathbb{Q}_p$  finite with ring of integers  $\mathcal{O} = \mathcal{O}_L$  and maximal ideal  $\lambda$ , and write  $\mathbb{F} = \mathcal{O}/\lambda$ . Let  $G$  be a profinite group which satisfies the following condition (Mazur's condition  $\Phi_p$ ): whenever  $\Delta$  is a finite index open subgroup of  $G$ , then  $\Delta/\langle[\Delta, \Delta], \Delta^p\rangle$  is finite. Equivalently (cf. Exercise 1.8.1 of [Boe]), for each  $\Delta$  the maximal pro- $p$  quotient of  $\Delta$  is topologically finitely generated. If  $G$  is topologically finitely generated, then  $\Phi_p$  holds, but we will need to use the condition for some  $G$  (the global Galois groups  $G_{K,S}$  defined below) which are not known to be topologically finitely generated.

In particular, using class field theory or Kummer theory, it can be checked that  $\Phi_p$  holds if  $G = G_K = \text{Gal}(\overline{K}/K)$  for some prime  $l$  (possibly equal to  $p$ ) and some finite extension  $K/\mathbb{Q}_l$ , or if  $G = G_{K,S} = \text{Gal}(K_S/K)$  where  $K$  is a number field,  $S$  is a finite set of finite places of  $K$ , and  $K_S/K$  is the maximal extension unramified outside of  $S$  and the infinite places (cf. the proof of Theorem 2.41 of [DDT97]).

Fix a representation  $\overline{\rho} : G \rightarrow \text{GL}_n(\mathbb{F})$ . Let  $\mathcal{C}_\mathcal{O}$  be the category of complete local noetherian  $\mathcal{O}$ -algebras with residue field  $\mathbb{F}$ , and consider the functor  $\mathcal{C}_\mathcal{O} \rightarrow \underline{\text{Sets}}$  which sends  $A$  to the set of continuous representations  $\rho : G \rightarrow \text{GL}_n(A)$  such that  $\rho \bmod \mathfrak{m}_A = \overline{\rho}$  (that is, to the set of lifts of  $\overline{\rho}$  to  $A$ ).

**3.2. Lemma.** *This functor is represented by a representation  $\rho^\square : G \rightarrow \text{GL}_n(R_{\overline{\rho}}^\square)$ .*

*Proof.* This is straightforward, cf. Proposition 1.3.1(a) of [Boe].  $\square$

**3.3. Definition.** We say that  $R_{\bar{\rho}}^{\square}$  is the *universal lifting ring* (or in Kisin's terminology, the *universal framed deformation ring*). We say that  $\rho^{\square}$  is the *universal lifting* of  $\bar{\rho}$ .

If  $\text{End}_{\mathbb{F}[G]}\bar{\rho} = \mathbb{F}$  we will say that  $\bar{\rho}$  is *Schur*. By Schur's lemma, if  $\bar{\rho}$  is absolutely irreducible, then  $\bar{\rho}$  is Schur. In this case, there is a very useful (and historically earlier) variant on the above construction.

**3.4. Definition.** Suppose that  $\bar{\rho}$  is Schur. Then a *deformation* of  $\bar{\rho}$  to  $A \in \text{ob } \mathcal{C}_{\mathcal{O}}$  is an equivalence class of liftings, where  $\rho \sim \rho'$  if and only if  $\rho' = a\rho a^{-1}$  for some  $a \in \ker(\text{GL}_n(A) \rightarrow \text{GL}_n(\mathbb{F}))$  (or equivalently, for some  $a \in \text{GL}_n(A)$ ).

**3.5. Lemma.** *If  $\bar{\rho}$  is Schur, then the functor  $\mathcal{C}_{\mathcal{O}} \rightarrow \underline{\text{Sets}}$  sending  $A$  to the set of deformations of  $\bar{\rho}$  to  $A$  is representable by some  $\rho^{\text{univ}} : G \rightarrow \text{GL}_n(R_{\bar{\rho}}^{\text{univ}})$ .*

*Proof.* See Proposition 1.3.1(b) of [Boe], or Theorem 2.36 of [DDT97] for a more hands-on approach.  $\square$

**3.6. Definition.** We say that  $\rho^{\text{univ}}$  (or more properly, its equivalence class) is the *universal deformation* of  $\bar{\rho}$ , and  $R_{\bar{\rho}}^{\text{univ}}$  is the *universal deformation ring*.

Deformations are representations considered up to conjugation, so it is reasonable to hope that deformations can be studied by considering their traces. In the case that  $\bar{\rho}$  is absolutely irreducible, universal deformations are determined by traces in the following rather strong sense. This result is essentially due to Carayol [Car94].

**3.7. Lemma.** *Suppose that  $\bar{\rho}$  is absolutely irreducible. Let  $R$  be an object of  $\mathcal{C}_{\mathcal{O}}$ , and  $\rho : G \rightarrow \text{GL}_n(R)$  a lifting of  $\bar{\rho}$ .*

- (1) *If  $a \in \text{GL}_n(R)$  and  $a\rho a^{-1} = \rho$  then  $a \in R^{\times}$ .*
- (2) *If  $\rho' : G \rightarrow \text{GL}_n(R)$  is another continuous lifting of  $\bar{\rho}$  and  $\text{tr } \rho = \text{tr } \rho'$ , then there is some  $a \in \ker(\text{GL}_n(R) \rightarrow \text{GL}_n(\mathbb{F}))$  such that  $\rho' = a\rho a^{-1}$ .*
- (3) *If  $S \subseteq R$  is a closed subring with  $S \in \text{ob } \mathcal{C}_{\mathcal{O}}$  and  $\mathfrak{m}_S = \mathfrak{m}_R \cap S$ , and if  $\text{tr } \rho(G) \subseteq S$ , then there is some  $a \in \ker(\text{GL}_n(R) \rightarrow \text{GL}_n(\mathbb{F}))$  such that  $a\rho a^{-1} : G \rightarrow \text{GL}_n(S)$ .*

*Proof.* See Lemmas 2.1.8 and 2.1.10 of [CHT08], or Theorem 2.2.1 of [Boe].  $\square$

**3.8. Exercise.** Deduce from Lemma 3.7 that if  $\bar{\rho}$  is absolutely irreducible, then  $R_{\bar{\rho}}^{\text{univ}}$  is topologically generated over  $\mathcal{O}$  by the values  $\text{tr } \rho^{\text{univ}}(g)$  as  $g$  runs over any dense subset of  $G$ .

**3.9. Exercise.** Show that if  $\bar{\rho}$  is absolutely irreducible, then  $R_{\bar{\rho}}^{\square}$  is isomorphic to a power series ring in  $(n^2 - 1)$  variables over  $R_{\bar{\rho}}^{\text{univ}}$ . Hint: let  $\rho^{\text{univ}}$  be a choice of universal deformation, and consider the homomorphism

$$\rho^{\square} : G \rightarrow \text{GL}_n(R_{\bar{\rho}}^{\text{univ}}[[X_{i,j}]_{i,j=1,\dots,n}/(X_{1,1})])$$

given by  $\rho^{\square} = (1_n + (X_{i,j}))\rho^{\text{univ}}(1_n + (X_{i,j}))^{-1}$ . Show that this is the universal lifting.

3.10. **Tangent spaces.** The tangent spaces of universal lifting and deformation rings have a natural interpretation in terms of liftings and deformations to the ring of dual numbers,  $\mathbb{F}[\varepsilon]/(\varepsilon^2)$ .

3.11. *Exercise.* Show that we have natural bijections between

- (1)  $\mathrm{Hom}_{\mathbb{F}}(\mathfrak{m}_{R_{\bar{\rho}}^{\square}} / (\mathfrak{m}_{R_{\bar{\rho}}^{\square}}^2, \lambda), \mathbb{F})$ .
- (2)  $\mathrm{Hom}_{\mathcal{O}}(R_{\bar{\rho}}^{\square}, \mathbb{F}[\varepsilon]/(\varepsilon^2))$ .
- (3) The set of liftings of  $\bar{\rho}$  to  $\mathbb{F}[\varepsilon]/(\varepsilon^2)$ .
- (4) The set of cocycles  $Z^1(G, \mathrm{ad} \bar{\rho})$ .

Show that if  $\bar{\rho}$  is absolutely irreducible, then we also have a bijection between  $\mathrm{Hom}_{\mathbb{F}}(\mathfrak{m}_{R_{\bar{\rho}}^{\mathrm{univ}}} / (\mathfrak{m}_{R_{\bar{\rho}}^{\mathrm{univ}}}^2, \lambda), \mathbb{F})$  and  $H^1(G, \mathrm{ad} \bar{\rho})$ . Hint: given  $f \in \mathrm{Hom}_{\mathbb{F}}(\mathfrak{m}_{R_{\bar{\rho}}^{\square}} / (\mathfrak{m}_{R_{\bar{\rho}}^{\square}}^2, \lambda), \mathbb{F})$ , define an element of  $\mathrm{Hom}_{\mathcal{O}}(R_{\bar{\rho}}^{\square}, \mathbb{F}[\varepsilon]/(\varepsilon^2))$  by sending  $a + x$  to  $a + f(x)\varepsilon$  whenever  $a \in \mathcal{O}$  and  $x \in \mathfrak{m}_{R_{\bar{\rho}}^{\square}}$ . Given a cocycle  $\phi \in Z^1(G, \mathrm{ad} \bar{\rho})$ , define a lifting  $\rho : G \rightarrow \mathrm{GL}_n(\mathbb{F}[\varepsilon]/(\varepsilon^2))$  by  $\rho(g) := (1 + \phi(g)\varepsilon)\bar{\rho}(g)$ .

3.12. **Corollary.** We have  $\dim_{\mathbb{F}} \mathfrak{m}_{R_{\bar{\rho}}^{\square}} / (\mathfrak{m}_{R_{\bar{\rho}}^{\square}}^2, \lambda) = \dim_{\mathbb{F}} H^1(G, \mathrm{ad} \bar{\rho}) + n^2 - \dim_{\mathbb{F}} H^0(G, \mathrm{ad} \bar{\rho})$ .

*Proof.* This follows from the exact sequence

$$0 \rightarrow (\mathrm{ad} \bar{\rho})^G \rightarrow \mathrm{ad} \bar{\rho} \rightarrow Z^1(G, \mathrm{ad} \bar{\rho}) \rightarrow H^1(G, \mathrm{ad} \bar{\rho}) \rightarrow 0. \quad \square$$

In particular, if  $d := \dim_{\mathbb{F}} Z^1(G, \mathrm{ad} \bar{\rho})$ , then we can choose a surjection  $\phi : \mathcal{O}[[x_1, \dots, x_d]] \twoheadrightarrow R_{\bar{\rho}}^{\square}$ . Similarly, if  $\bar{\rho}$  is absolutely irreducible, we can choose a surjection  $\phi' : \mathcal{O}[[x_1, \dots, x_{d'}]] \twoheadrightarrow R_{\bar{\rho}}^{\mathrm{univ}}$ , where  $d' := \dim_{\mathbb{F}} H^1(G, \mathrm{ad} \bar{\rho})$ .

3.13. **Lemma.** If  $J = \ker \phi$  or  $J = \ker \phi'$ , then there is an injection  $\mathrm{Hom}_{\mathbb{F}}(J/\mathfrak{m}J, \mathbb{F}) \hookrightarrow H^2(G, \mathrm{ad} \bar{\rho})$ .

*Proof.* See Proposition 2 of [Maz89]. □

3.14. **Corollary.** If  $H^2(G, \mathrm{ad} \bar{\rho}) = (0)$ , then  $R_{\bar{\rho}}^{\square}$  is formally smooth of relative dimension  $\dim_{\mathbb{F}} Z^1(G, \mathrm{ad} \bar{\rho})$  over  $\mathcal{O}$ .

In any case, the Krull dimension of  $R_{\bar{\rho}}^{\square}$  is at least

$$1 + n^2 - \dim_{\mathbb{F}} H^0(G, \mathrm{ad} \bar{\rho}) + \dim_{\mathbb{F}} H^1(G, \mathrm{ad} \bar{\rho}) - \dim_{\mathbb{F}} H^2(G, \mathrm{ad} \bar{\rho}).$$

If  $\bar{\rho}$  is absolutely irreducible, then the Krull dimension of  $R_{\bar{\rho}}^{\mathrm{univ}}$  is at least

$$1 + \dim_{\mathbb{F}} H^1(G, \mathrm{ad} \bar{\rho}) - \dim_{\mathbb{F}} H^2(G, \mathrm{ad} \bar{\rho}).$$

3.15. **Deformation conditions.** In practice, we frequently want to impose additional conditions on the liftings and deformations we consider. For example, if we are trying to prove the Fontaine–Mazur conjecture, we would like to be able to restrict to global deformations which are geometric. There are various ways in which to impose extra conditions; we will use the formalism of *deformation problems* introduced in [CHT08].

**3.16. Definition.** By a *deformation problem*  $\mathcal{D}$  we mean a collection of liftings  $(R, \rho)$  of  $(\mathbb{F}, \bar{\rho})$  (with  $R$  an object of  $\mathcal{C}_{\mathcal{O}}$ ), satisfying the following properties.

- $(\mathbb{F}, \bar{\rho}) \in \mathcal{D}$ .
- If  $f : R \rightarrow S$  is a morphism in  $\mathcal{C}_{\mathcal{O}}$  and  $(R, \rho) \in \mathcal{D}$ , then  $(S, f \circ \rho) \in \mathcal{D}$ .
- If  $f : R \hookrightarrow S$  is a morphism in  $\mathcal{C}_{\mathcal{O}}$  then  $(R, \rho) \in \mathcal{D}$  if and only if  $(S, f \circ \rho) \in \mathcal{D}$ .
- Suppose that  $R_1, R_2 \in \text{ob } \mathcal{C}_{\mathcal{O}}$  and  $I_1, I_2$  are ideals of  $R_1, R_2$  respectively such that there is an isomorphism  $f : R_1/I_1 \xrightarrow{\sim} R_2/I_2$ . Suppose also that  $(R_1, \rho_1), (R_2, \rho_2) \in \mathcal{D}$ , and that  $f(\rho_1 \bmod I_1) = \rho_2 \bmod I_2$ .  
Then  $(\{(a, b) \in R_1 \oplus R_2 : f(a \bmod I_1) = b \bmod I_2\}, \rho_1 \oplus \rho_2) \in \mathcal{D}$ .
- If  $(R, \rho)$  is a lifting of  $(\mathbb{F}, \bar{\rho})$  and  $I_1 \supset I_2 \supset \cdots$  is a sequence of ideals of  $R$  with  $\bigcap_j I_j = (0)$ , and  $(R/I_j, \rho \bmod I_j) \in \mathcal{D}$  for all  $j$ , then  $(R, \rho) \in \mathcal{D}$ .
- If  $(R, \rho) \in \mathcal{D}$  and  $a \in \ker(\text{GL}_n(R) \rightarrow \text{GL}_n(\mathbb{F}))$ , then  $(R, a\rho a^{-1}) \in \mathcal{D}$ .

In practice, when we want to impose a condition on our deformations, it will be easy to see that it satisfies these requirements. (An exception is that these properties are hard to check for certain conditions arising in  $p$ -adic Hodge theory, but we won't need those conditions in this course.)

The relationship of this definition to the universal lifting ring is as follows. Note that each element  $a \in \ker(\text{GL}_n(R_{\bar{\rho}}^{\square}) \rightarrow \text{GL}_n(\mathbb{F}))$  acts on  $R_{\bar{\rho}}^{\square}$ , via the universal property and by sending  $\rho^{\square}$  to  $a^{-1}\rho^{\square}a$ . [Warning: this *isn't* a group action, though!]

**3.17. Lemma.** (1) If  $\mathcal{D}$  is a deformation problem then there is a  $\ker(\text{GL}_n(R_{\bar{\rho}}^{\square}) \rightarrow \text{GL}_n(\mathbb{F}))$ -invariant ideal  $I(\mathcal{D})$  of  $R_{\bar{\rho}}^{\square}$  such that  $(R, \rho) \in \mathcal{D}$  if and only if the map  $R_{\bar{\rho}}^{\square} \rightarrow R$  induced by  $\rho$  factors through the quotient  $R_{\bar{\rho}}^{\square}/I(\mathcal{D})$ .

(2) Let  $\tilde{L}(\mathcal{D}) \subseteq Z^1(G, \text{ad } \bar{\rho}) \cong \text{Hom}(m_{R_{\bar{\rho}}^{\square}}/(\lambda, m_{R_{\bar{\rho}}^{\square}}^2), \mathbb{F})$  denote the annihilator of the image of  $I(\mathcal{D})$  in  $m_{R_{\bar{\rho}}^{\square}}/(\lambda, m_{R_{\bar{\rho}}^{\square}}^2)$ .

Then  $\tilde{L}(\mathcal{D})$  is the preimage of some subspace  $L(\mathcal{D}) \subseteq H^1(G, \text{ad } \bar{\rho})$ .

(3) If  $I$  is a  $\ker(\text{GL}_n(R_{\bar{\rho}}^{\square}) \rightarrow \text{GL}_n(\mathbb{F}))$ -invariant ideal of  $R_{\bar{\rho}}^{\square}$  with  $\sqrt{I} = I$  and  $I \neq m_{R_{\bar{\rho}}^{\square}}$ , then

$$\mathcal{D}(I) := \{(R, \rho) : R_{\bar{\rho}}^{\square} \rightarrow R \text{ factors through } R_{\bar{\rho}}^{\square}/I\}$$

is a deformation problem. Furthermore, we have  $I(\mathcal{D}(I)) = I$  and  $\mathcal{D}(I(\mathcal{D})) = \mathcal{D}$ .

*Proof.* See Lemma 2.2.3 of [CHT08] and Lemma 3.2 of [BLGHT09] (and for (2), use that  $I(\mathcal{D})$  is  $\ker(\text{GL}_n(R_{\bar{\rho}}^{\square}) \rightarrow \text{GL}_n(\mathbb{F}))$ -invariant).  $\square$

**3.18. Fixing determinants.** For technical reasons, we will want to fix the determinants of our Galois representations. To this end, let  $\chi : G \rightarrow \mathcal{O}^{\times}$  be a continuous homomorphism such that  $\chi \bmod \lambda = \det \bar{\rho}$ . Then it makes sense to ask that a lifting has determinant  $\chi$ , and we



can define a universal lifting ring  $R_{\bar{\rho}, \chi}^{\square}$  for lifts with determinant  $\chi$ , and when  $\bar{\rho}$  is Schur, a universal fixed determinant deformation ring  $R_{\bar{\rho}, \chi}^{\text{univ}}$ .

3.19. *Exercise.* Check that the material developed in the previous section goes over unchanged, except that  $\text{ad } \bar{\rho}$  needs to be replaced with  $\text{ad } {}^0\bar{\rho} := \{x \in \text{ad } \bar{\rho} : \text{tr } x = 0\}$ .

Note that since we are assuming throughout that  $p \nmid n$ ,  $\text{ad } {}^0\bar{\rho}$  is a direct summand of  $\text{ad } \bar{\rho}$  (as a  $G$ -representation).

3.20. **Global deformations with local conditions.** Now fix a finite set  $S$ , and for each  $v \in S$ , a profinite group  $G_v$  satisfying  $\Phi_p$ , together with a continuous homomorphism  $G_v \rightarrow G$ , and a deformation problem  $\mathcal{D}_v$  for  $\bar{\rho}|_{G_v}$ . [In applications,  $G$  will be a global Galois group, and the  $G_v$  will be decomposition groups at finite places.]

Also fix  $\chi : G \rightarrow \mathcal{O}^{\times}$ , a continuous homomorphism such that  $\chi \bmod \lambda = \det \bar{\rho}$ . Assume that  $\bar{\rho}$  is absolutely irreducible, and fix some subset  $T \subseteq S$ .

3.21. **Definition.** Fix  $A \in \text{ob } \mathcal{C}_{\mathcal{O}}$ . A  $T$ -framed deformation of  $\bar{\rho}$  of type  $\mathcal{S} := (S, \{\mathcal{D}_v\}_{v \in S}, \chi)$  to  $A$  is an equivalence class of tuples  $(\rho, \{\alpha_v\}_{v \in T})$ , where  $\rho : G \rightarrow \text{GL}_n(A)$  is a lift of  $\bar{\rho}$  such that  $\det \rho = \chi$  and  $\rho|_{G_v} \in \mathcal{D}_v$  for all  $v \in S$ , and  $\alpha_v$  is an element of  $\ker(\text{GL}_n(A) \rightarrow \text{GL}_n(\mathbb{F}))$ .

The equivalence relation is defined by decreeing that for each  $\beta \in \ker(\text{GL}_n(A) \rightarrow \text{GL}_n(\mathbb{F}))$ , we have  $(\rho, \{\alpha_v\}_{v \in T}) \sim (\beta\rho\beta^{-1}, \{\beta\alpha_v\}_{v \in T})$ .

The point of considering  $T$ -framed deformations is that it allows us to study absolutely irreducible representations  $\bar{\rho}$  for which some of the  $\bar{\rho}|_{G_v}$  are reducible, because if  $(\rho, \{\alpha_v\}_{v \in T})$  is a  $T$ -framed deformation of type  $\mathcal{S}$ , then  $\alpha_v^{-1}\rho|_{G_v}\alpha_v$  is a well-defined element of  $\mathcal{D}_v$  (independent of the choice of representative of the equivalence class). The following lemma should be unsurprising.

3.22. **Lemma.** *The functor  $\mathcal{C}_{\mathcal{O}} \rightarrow \text{Sets}$  sending  $A$  to the set of  $T$ -framed deformations of  $\bar{\rho}$  of type  $\mathcal{S}$  is represented by a universal object  $\rho^{\square T} : G \rightarrow \text{GL}_n(R_S^{\square T})$ .*

*Proof.* See Proposition 2.2.9 of [CHT08]. □

If  $T = \emptyset$  then we will write  $R_S^{\text{univ}}$  for  $R_S^{\square T}$ .

3.23. **Presenting global deformation rings over local lifting rings.** Continue to use the notation of the previous subsection. Since  $\alpha_v^{-1}\rho^{\square T}|_{G_v}\alpha_v$  is a well-defined element of  $\mathcal{D}_v$ , we have a tautological homomorphism  $R_{\bar{\rho}|_{G_v}, \chi}^{\square} / I(\mathcal{D}_v) \rightarrow R_S^{\square T}$ . Define

$$R_{S, T}^{\text{loc}} := \widehat{\otimes}_{v \in T} \left( R_{\bar{\rho}|_{G_v}, \chi}^{\square} / I(\mathcal{D}_v) \right).$$

Then we have a natural map  $R_{S, T}^{\text{loc}} \rightarrow R_S^{\square T}$ .

We now generalise Corollary 3.14 by considering presentations of  $R_S^{\square T}$  over  $R_{S,T}^{\text{loc}}$ . In order to compute how many variables are needed to present  $R_S^{\square T}$  over  $R_{S,T}^{\text{loc}}$ , we must compute  $\dim_{\mathbb{F}} \mathfrak{m}_{R_S^{\square T}} / (\mathfrak{m}_{R_S^{\square T}}^2, \mathfrak{m}_{R_{S,T}^{\text{loc}}}, \lambda)$ . Unsurprisingly, in order to compute this, we will compute a certain  $H^1$ .

We define a complex as follows. As usual, given a group  $G$  and a  $\mathbb{F}[G]$ -module  $M$ , we let  $C^i(G, M)$  be the space of functions  $G^i \rightarrow M$ , and we let  $\partial : C^i(G, M) \rightarrow C^{i+1}(G, M)$  be the coboundary map. We define a complex  $C_{S,T,\text{loc}}^i(G, \text{ad } {}^0\bar{\rho})$  by

$$\begin{aligned} C_{S,T,\text{loc}}^0(G, \text{ad } {}^0\bar{\rho}) &= \bigoplus_{v \in T} C^0(G_v, \text{ad } \bar{\rho}) \oplus \bigoplus_{v \in S \setminus T} 0, \\ C_{S,T,\text{loc}}^1(G, \text{ad } {}^0\bar{\rho}) &= \bigoplus_{v \in T} C^1(G_v, \text{ad } {}^0\bar{\rho}) \oplus \bigoplus_{v \in S \setminus T} C^1(G_v, \text{ad } {}^0\bar{\rho}) / \tilde{L}(\mathcal{D}_v), \end{aligned}$$

and for  $i \geq 2$ ,

$$C_{S,T,\text{loc}}^i(G, \text{ad } {}^0\bar{\rho}) = \bigoplus_{v \in S} C^i(G_v, \text{ad } {}^0\bar{\rho}).$$

Let  $C_0^0(G, \text{ad } {}^0\bar{\rho}) := C^0(G, \text{ad } \bar{\rho})$ , and set  $C_0^i(G, \text{ad } {}^0\bar{\rho}) = C^i(G, \text{ad } {}^0\bar{\rho})$  for  $i > 0$ . Then we let  $H_{S,T}^i(G, \text{ad } {}^0\bar{\rho})$  denote the cohomology of the complex

$$C_{S,T}^i(G, \text{ad } {}^0\bar{\rho}) := C_0^i(G, \text{ad } {}^0\bar{\rho}) \oplus C_{S,T,\text{loc}}^{i-1}(G, \text{ad } {}^0\bar{\rho})$$

where the coboundary map is given by

$$(\phi, (\psi_v)) \mapsto (\partial\phi, (\phi|_{G_v} - \partial\psi_v)).$$

Then we have an exact sequence of complexes

$$0 \rightarrow C_{S,T,\text{loc}}^{i-1}(G, \text{ad } {}^0\bar{\rho}) \rightarrow C_{S,T}^i(G, \text{ad } {}^0\bar{\rho}) \rightarrow C_0^i(G, \text{ad } {}^0\bar{\rho}) \rightarrow 0,$$

and the corresponding long exact sequence in cohomology is

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_{S,T}^0(G, \text{ad } {}^0\bar{\rho}) & \longrightarrow & H^0(G, \text{ad } \bar{\rho}) & \longrightarrow & \bigoplus_{v \in T} H^0(G_v, \text{ad } \bar{\rho}) \\ & & \longleftarrow & & \longleftarrow & & \longleftarrow \\ & & H_{S,T}^1(G, \text{ad } {}^0\bar{\rho}) & \longrightarrow & H^1(G, \text{ad } {}^0\bar{\rho}) & \longrightarrow & \bigoplus_{v \in T} H^1(G_v, \text{ad } {}^0\bar{\rho}) \oplus \bigoplus_{v \in S \setminus T} H^1(G_v, \text{ad } {}^0\bar{\rho}) / L(\mathcal{D}_v) \\ & & \longleftarrow & & \longleftarrow & & \longleftarrow \\ & & H_{S,T}^2(G, \text{ad } {}^0\bar{\rho}) & \longrightarrow & H^2(G, \text{ad } {}^0\bar{\rho}) & \longrightarrow & \bigoplus_{v \in S} H^2(G_v, \text{ad } {}^0\bar{\rho}) \\ & & \longleftarrow & & \longleftarrow & & \longleftarrow \\ & & H_{S,T}^3(G, \text{ad } {}^0\bar{\rho}) & \longrightarrow & \dots & & \end{array}$$

Taking Euler characteristics, we see that if we define e.g.  $\chi(G, \text{ad } {}^0\bar{\rho}) = \sum_i (-1)^{i-1} \dim_{\mathbb{F}} H^i(G, \text{ad } {}^0\bar{\rho})$ , we have

$$\begin{aligned} \chi_{S,T}(G, \text{ad } {}^0\bar{\rho}) &= -1 + \chi(G, \text{ad } {}^0\bar{\rho}) - \sum_{v \in S} \chi(G_v, \text{ad } {}^0\bar{\rho}) + \sum_{v \in T} (\dim_{\mathbb{F}} H^0(G_v, \text{ad } \bar{\rho}) - \dim_{\mathbb{F}} H^0(G_v, \text{ad } {}^0\bar{\rho})) \\ &\quad + \sum_{v \in S \setminus T} (\dim_{\mathbb{F}} L(\mathcal{D}_v) - \dim_{\mathbb{F}} H^0(G_v, \text{ad } {}^0\bar{\rho})). \end{aligned}$$

From now on for the rest of the notes, we specialise to the case that  $F$  is a number field,  $S$  is a finite set of finite places of  $F$  including all the places lying over  $p$ , and we set  $G = G_{F,S}$ ,  $G_v =$

$G_{F_v}$  for  $v \in S$ . (Since  $G = G_{F,S}$ , note in particular that all deformations we are considering are unramified outside of  $S$ .) We then employ standard results on Galois cohomology that can be found in [Mil06]. In particular, we have  $H^n(G_{F_v}, \text{ad } \bar{\rho}) = (0)$  if  $n \geq 3$ , and

$$H^n(G_{F,S}, \text{ad } \bar{\rho}) \cong \bigoplus_{v \text{ real}} H^n(G_{F_v}, \text{ad } \bar{\rho}) = (0)$$

if  $n \geq 3$  (the vanishing of the local cohomology groups follows as  $p > 2$ , so  $G_{F_v}$  has order coprime to that of  $\text{ad } \bar{\rho}$ ). Consequently,  $H_{S,T}^n(G_{F,S}, \text{ad } \bar{\rho}) = (0)$  if  $n > 3$ .

We now employ the local and global Euler characteristic formulas. For simplicity, assume from now on that  $T$  contains all the places of  $S$  lying over  $p$ . The global formula gives

$$\chi(G_{F,S}, \text{ad } \bar{\rho}) = - \sum_{v|\infty} \dim_{\mathbb{F}} H^0(G_{F_v}, \text{ad } \bar{\rho}) + [F : \mathbb{Q}](n^2 - 1),$$

and the local formula gives

$$\begin{aligned} \sum_{v \in S} \chi(G_{F_v}, \text{ad } \bar{\rho}) &= \sum_{v|p} (n^2 - 1)[F_v : \mathbb{Q}_p] \\ &= (n^2 - 1)[F : \mathbb{Q}], \end{aligned}$$

so that

$$\chi_{S,T}(G_{F,S}, \text{ad } \bar{\rho}) = -1 + \#T - \sum_{v|\infty} \dim_{\mathbb{F}} H^0(G_{F_v}, \text{ad } \bar{\rho}) + \sum_{v \in S \setminus T} \left( \dim_{\mathbb{F}} L(\mathcal{D}_v) - \dim_{\mathbb{F}} H^0(G_{F_v}, \text{ad } \bar{\rho}) \right).$$

Assume now that  $\bar{\rho}$  is absolutely irreducible; then  $H^0(G_{F,S}, \text{ad } \bar{\rho}) = \mathbb{F}$ , so  $H_{S,T}^0(G_{F,S}, \text{ad } \bar{\rho}) = \mathbb{F}$ . To say something sensible about  $H_{S,T}^1(G_{F,S}, \text{ad } \bar{\rho})$  we still need to control the  $H_{S,T}^2$  and  $H_{S,T}^3$ . Firstly, the above long exact sequence gives us in particular the exact sequence

$$\begin{array}{ccccccc} & & H^1(G_{F,S}, \text{ad } \bar{\rho}) & \longrightarrow & \bigoplus_{v \in T} H^1(G_{F_v}, \text{ad } \bar{\rho}) \oplus \bigoplus_{v \in S \setminus T} H^1(G_{F_v}, \text{ad } \bar{\rho}) / L(\mathcal{D}_v) & & \\ \longleftarrow & & & & & & \\ & \longrightarrow & H_{S,T}^2(G_{F,S}, \text{ad } \bar{\rho}) & \longrightarrow & H^2(G_{F,S}, \text{ad } \bar{\rho}) & \longrightarrow & \bigoplus_{v \in S} H^2(G_{F_v}, \text{ad } \bar{\rho}) \\ \longleftarrow & & & & & & \\ & \longrightarrow & H_{S,T}^3(G_{F,S}, \text{ad } \bar{\rho}) & \longrightarrow & 0. & & \end{array}$$

On the other hand, from the Poitou–Tate exact sequence [Mil06, Prop. 4.10, Chapter 1] we have an exact sequence

$$\begin{array}{ccccccc} & & H^1(G_{F,S}, \text{ad } \bar{\rho}) & \longrightarrow & \bigoplus_{v \in S} H^1(G_{F_v}, \text{ad } \bar{\rho}) & \longrightarrow & H^1(G_{F,S}, (\text{ad } \bar{\rho})^\vee(1))^\vee \\ \longleftarrow & & & & & & \\ & \longrightarrow & H^2(G_{F,S}, \text{ad } \bar{\rho}) & \longrightarrow & \bigoplus_{v \in S} H^2(G_{F_v}, \text{ad } \bar{\rho}) & \longrightarrow & H^0(G_{F,S}, (\text{ad } \bar{\rho})^\vee(1))^\vee \longrightarrow 0. \end{array}$$

Note that  $\text{ad } \bar{\rho}$  is self-dual under the trace pairing, so we can and do identify  $(\text{ad } \bar{\rho})^\vee(1)$  and  $(\text{ad } \bar{\rho})(1)$ . If we let  $L(\mathcal{D}_v)^\perp \subseteq H^1(G_{F_v}, (\text{ad } \bar{\rho})(1))$  denote the annihilator of  $L(\mathcal{D}_v)$  under the pairing coming from Tate local duality, and we define

$$H_{S,T}^1(G_{F,S}, (\text{ad } \bar{\rho})(1)) := \ker \left( H^1(G_{F,S}, (\text{ad } \bar{\rho})(1)) \rightarrow \bigoplus_{v \in S \setminus T} \left( H^1(G_{F_v}, (\text{ad } \bar{\rho})(1)) / L(\mathcal{D}_v)^\perp \right) \right),$$

then we deduce that we have an exact sequence

$$\begin{array}{ccccccc}
& & H^1(G_{F,S}, \text{ad } {}^0\bar{\rho}) & \longrightarrow & \oplus_{v \in T} H^1(G_{F_v}, \text{ad } {}^0\bar{\rho}) \oplus_{v \in S \setminus T} H^1(G_{F_v}, \text{ad } {}^0\bar{\rho}) / L(\mathcal{D}_v) & & \\
\longleftarrow & & & & & & \longrightarrow \\
& & H^1_{S,T}(G_{F,S}, \text{ad } {}^0\bar{\rho}(1))^\vee & \longrightarrow & H^2(G_{F,S}, \text{ad } {}^0\bar{\rho}) & \longrightarrow & \oplus_{v \in S} H^2(G_{F_v}, \text{ad } {}^0\bar{\rho}) \\
& & & & & & \longleftarrow \\
& & H^0(G_{F,S}, \text{ad } {}^0\bar{\rho}(1))^\vee & \longrightarrow & 0, & & 
\end{array}$$

and comparing with the diagram above shows that

$$\begin{aligned}
H^3_{S,T}(G_{F,S}, \text{ad } {}^0\bar{\rho}) &\cong H^0(G_{F,S}, \text{ad } {}^0\bar{\rho}(1))^\vee, \\
H^2_{S,T}(G_{F,S}, \text{ad } {}^0\bar{\rho}) &\cong H^1_{S,T}(G_{F,S}, \text{ad } {}^0\bar{\rho}(1))^\vee.
\end{aligned}$$

Combining all of this, we see that

$$\begin{aligned}
\dim_{\mathbb{F}} H^1_{S,T}(G_{F,S}, \text{ad } {}^0\bar{\rho}) &= \#T - \sum_{v|\infty} \dim_{\mathbb{F}} H^0(G_{F_v}, \text{ad } {}^0\bar{\rho}) + \sum_{v \in S \setminus T} \left( \dim_{\mathbb{F}} L(\mathcal{D}_v) - \dim_{\mathbb{F}} H^0(G_{F_v}, \text{ad } {}^0\bar{\rho}) \right) \\
&\quad + \dim_{\mathbb{F}} H^1_{S,T}(G_{F,S}, \text{ad } {}^0\bar{\rho}(1)) - \dim_{\mathbb{F}} H^0(G_{F,S}, \text{ad } {}^0\bar{\rho}(1)).
\end{aligned}$$

Now, similar arguments to those we used above give us the following result (cf. Section 2.2 of [CHT08]).

**3.24. Proposition.** (1) *There is a canonical isomorphism*

$$\text{Hom}(\mathfrak{m}_{R_S^{\square T}} / (\mathfrak{m}_{R_S^{\square T}}^2, \mathfrak{m}_{R_S^{\text{loc}}}, \lambda), \mathbb{F}) \cong H^1_{S,T}(G_{F,S}, \text{ad } {}^0\bar{\rho}).$$

(2)  $R_S^{\square T}$  is the quotient of a power series ring in  $\dim_{\mathbb{F}} H^1_{S,T}(G_{F,S}, \text{ad } {}^0\bar{\rho})$  variables over  $R_S^{\text{loc}}$ .

(3) The Krull dimension of  $R_S^{\text{univ}}$  is at least

$$1 + \sum_{v \in S} \left( \text{Krull dim.}(R_{\bar{\rho}|_{G_{F_v}, \chi}}^{\square} / I(\mathcal{D}_v)) - n^2 \right) - \sum_{v|\infty} \dim_{\mathbb{F}} H^0(G_{F_v}, \text{ad } {}^0\bar{\rho}) - \dim_{\mathbb{F}} H^0(G_{F,S}, \text{ad } {}^0\bar{\rho}(1)).$$

**3.25. Finiteness of maps between global deformation rings.** Suppose that  $F'/F$  is a finite extension of number fields, and that  $S'$  is the set of places of  $F'$  lying over  $S$ . Assume that  $\bar{\rho}|_{G_{F', S'}}$  is absolutely irreducible. Then restricting the universal deformation  $\rho^{\text{univ}}$  of  $\bar{\rho}$  to  $G_{F', S'}$  gives a ring homomorphism  $R_{\bar{\rho}|_{G_{F', S'}}}^{\text{univ}} \rightarrow R_{\bar{\rho}}^{\text{univ}}$ . The following very useful fact is due to Khare and Wintenberger.

**3.26. Proposition.** *The ring  $R_{\bar{\rho}}^{\text{univ}}$  is a finitely generated  $R_{\bar{\rho}|_{G_{F', S'}}}^{\text{univ}}$ -module.*

*Proof.* See e.g. Lemma 1.2.3 of [BLGGT10]. (Note that it is easy to see that it is quasifinite, but finiteness uses Exercise 3.8.)  $\square$

**3.27. Local deformation rings with  $l = p$ .** The local deformation rings when  $l = p$  are one of the most difficult and interesting parts of the subject; for example, a detailed computation of deformation rings with  $l = p = 3$  was at the heart of the eventual proof of the Taniyama–Shimura–Weil conjecture, and many of the recent improvements to automorphy lifting theorems have relied on deep results on these deformation rings due to Kisin.

In this course, we will ignore all of these difficulties, and work only with the “Fontaine–Laffaille” case. This is already enough to have important applications. In this case the local deformation rings are smooth; for the most part, the deformation rings that we care about when  $l = p$  aren’t smooth, but the situation is in some sense worse than this, in that we don’t have a concrete description of the rings in most cases, or even basic information such as the number of irreducible components of the generic fibre.

Assume that  $K/\mathbb{Q}_p$  is a finite unramified extension, and assume that  $L$  is chosen large enough to contain the images of all embeddings  $K \hookrightarrow \overline{\mathbb{Q}_p}$ . For each  $\sigma : K \hookrightarrow L$ , let  $H_\sigma$  be a set of  $n$  distinct integers, such that the difference between the maximal and minimal elements of  $H_\sigma$  is less than or equal to  $p - 2$ .

**3.28. Theorem.** *There is a unique reduced,  $p$ -torsion free quotient  $R_{\rho, \chi, \text{cr}, \{H_\sigma\}}^\square$  of  $R_{\rho, \chi}^\square$  with the property that a continuous homomorphism  $\psi : R_{\rho, \chi}^\square \rightarrow \overline{\mathbb{Q}_p}$  factors through  $R_{\rho, \chi, \text{cr}, \{H_\sigma\}}^\square$  if and only if  $\psi \circ \rho^\square$  is crystalline, and for each  $\sigma : K \hookrightarrow L$ , we have  $\text{HT}_\sigma(\psi \circ \rho^\square) = H_\sigma$ .*

*Furthermore,  $\dim R_{\rho, \chi, \text{cr}, \{H_\sigma\}}^\square = n^2 + [F_v : \mathbb{Q}_p] \frac{1}{2}n(n-1)$ , and in fact  $R_{\rho, \chi, \text{cr}, \{H_\sigma\}}^\square$  is formally smooth over  $\mathcal{O}$ , i.e. it is isomorphic to a power series ring in  $n^2 - 1 + [F_v : \mathbb{Q}_p] \frac{1}{2}n(n-1)$  variables over  $\mathcal{O}$ .*

In fact, if we remove the assertion of formal smoothness, Theorem 3.28 still holds without the assumption that  $K/\mathbb{Q}_p$  is unramified, and without any assumption on the difference between the maximal and minimal elements of the  $H_\sigma$ , but in this case it is a much harder theorem of Kisin ([Kis08]). In any case, the formal smoothness will be important for us.

Theorem 3.28 is essentially a consequence of Fontaine–Laffaille theory [FL82], which is a form of integral  $p$ -adic Hodge theory; that is, it classifies the Galois-stable lattices in crystalline representations, under the assumptions we’ve made above. The first proof of Theorem 3.28 was essentially in Ramakrishna’s thesis [Ram93], and the general result is the content of section 2.4 of [CHT08].

**3.29. Local deformation rings with  $p \neq l$ .** In contrast to the situation when  $l = p$ , we will need to consider several deformation problems when  $l \neq p$ . We will restrict ourselves to the two-dimensional case. Let  $K/\mathbb{Q}_l$  be a finite extension, with  $l \neq p$ , and fix  $n = 2$ . As we saw in Section 2.6, there is essentially an incompatibility between the wild inertia subgroup of  $G_K$  and the  $p$ -adic topology on  $\text{GL}_2(\mathcal{O})$ , which makes it possible to explicitly describe the  $p$ -adic representations of  $G_K$ , and consequently the corresponding universal deformation rings. This was done in varying degrees of generality over a long period of time; in particular, in

the general  $n$ -dimensional case we highlight Section 2.4.4 of [CHT08] and [Cho09], and in the 2-dimensional setting [Pil] and [Sho13]. In fact [Sho13] gives a complete description of the deformation rings for a fixed inertial type.

We will content ourselves with recalling some of the basic structural results, and with giving a sketch of how the results are proved in one particular case (see Exercise 3.34 below).

**3.30. Deformations of fixed type.** Recall from Proposition 2.17 that given a representation  $\rho : G_K \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}}_p)$  there is a Weil–Deligne representation  $\mathrm{WD}(\rho)$  associated to  $\rho$ . If  $\mathrm{WD} = (r, N)$  is a Weil–Deligne representation, then we write  $\mathrm{WD}|_{I_K}$  for  $(r|_{I_K}, N)$ , and call it an *inertial WD-type*.

Fix  $\bar{\rho} : G_K \rightarrow \mathrm{GL}_2(\mathbb{F})$ . Then (assuming as usual that  $L$  is sufficiently large) we have the following general result on  $R_{\bar{\rho}, \chi}^{\square}$  (see e.g. Theorem 3.3.1 of [Boe]).

**3.31. Theorem.**  $R_{\bar{\rho}, \chi}^{\square}$  is equidimensional of Krull dimension 4, and the generic fibre  $R_{\bar{\rho}, \chi}^{\square}[1/p]$  has Krull dimension 3. Furthermore:

(a) The function which takes a  $\overline{\mathbb{Q}}_p$ -point  $x : R_{\bar{\rho}, \chi}^{\square}[1/p] \rightarrow \overline{\mathbb{Q}}_p$  to (the isomorphism class of)  $\mathrm{WD}(x \circ \rho^{\square})|_{I_K}$  (forgetting  $N$ ) is constant on the irreducible components of  $R_{\bar{\rho}, \chi}^{\square}[1/p]$ .

(b) The irreducible components of  $R_{\bar{\rho}, \chi}^{\square}[1/p]$  are all formally smooth, and there are only finitely many of them.

In the light of Theorem 3.31, we make the following definition. Let  $\tau$  be an inertial WD-type. Then there is a unique reduced,  $p$ -torsion free quotient  $R_{\bar{\rho}, \chi, \tau}^{\square}$  of  $R_{\bar{\rho}, \chi}^{\square}$  with the property that a continuous homomorphism  $\psi : R_{\bar{\rho}, \chi}^{\square} \rightarrow \overline{\mathbb{Q}}_p$  factors through  $R_{\bar{\rho}, \chi, \tau}^{\square}$  if and only if  $\psi \circ \rho^{\square}$  has inertial Weil–Deligne type  $\tau$ . (Of course, for all but finitely many  $\tau$ , we will just have  $R_{\bar{\rho}, \chi, \tau}^{\square} = 0$ .) By Theorem 3.31 we see that if  $R_{\bar{\rho}, \chi, \tau}^{\square}$  is nonzero then it has Krull dimension 4, and its generic fibre is irreducible and formally smooth.

**3.32. Taylor–Wiles deformations.** As the name suggests, the deformations that we consider in this subsection will be of crucial importance for the Taylor–Wiles–Kisin method. Write  $k$  for the residue field of  $K$ , and suppose that  $\bar{\rho}$  is unramified, that  $\bar{\rho}(\mathrm{Frob}_K)$  has distinct eigenvalues, and that  $\#k \equiv 1 \pmod{p}$ . Suppose also that  $\chi$  is unramified.

**3.33. Lemma.** Suppose that  $(\#k - 1)$  is exactly divisible by  $p^m$ . Then  $R_{\bar{\rho}, \chi}^{\square} \cong \mathcal{O}[[x, y, B, u]] / ((1 + u)^{p^m} - 1)$ . Furthermore, if  $\varphi \in G_K$  is a lift of  $\mathrm{Frob}_K$ , then  $\rho^{\square}(\varphi)$  is conjugate to a diagonal matrix.

**3.34. Exercise.** Prove this lemma as follows. Note firstly that  $\rho^{\square}(P_K) = \{1\}$ , because  $\bar{\rho}(P_K) = \{1\}$ , so  $\rho^{\square}(P_K)$  is a pro- $l$ -subgroup of the pro- $p$ -group  $\ker(\mathrm{GL}_2(R_{\bar{\rho}, \chi}^{\square}) \rightarrow \mathrm{GL}_2(\mathbb{F}))$ .

Let  $\varphi$  be a fixed lift of  $\mathrm{Frob}_K$  to  $G_K/P_K$ , and  $\sigma$  a topological generator of  $I_K/P_K$ , which as in Section 2.6 we can choose so that  $\varphi^{-1}\sigma\varphi = \sigma^{\#k}$ . Write  $\bar{\rho}(\varphi) = \begin{pmatrix} \bar{\alpha} & 0 \\ 0 & \bar{\beta} \end{pmatrix}$ , and fix lifts  $\alpha, \beta \in \mathcal{O}$  of  $\bar{\alpha}, \bar{\beta}$ .

Then we will show that we can take

$$\rho^\square(\varphi) = \begin{pmatrix} 1 & y \\ x & 1 \end{pmatrix}^{-1} \begin{pmatrix} \alpha + B & 0 \\ 0 & \chi(\varphi)/(\alpha + B) \end{pmatrix} \begin{pmatrix} 1 & y \\ x & 1 \end{pmatrix},$$

$$\rho^\square(\sigma) = \begin{pmatrix} 1 & y \\ x & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 + u & 0 \\ 0 & (1 + u)^{-1} \end{pmatrix} \begin{pmatrix} 1 & y \\ x & 1 \end{pmatrix}.$$

- (1) Let  $\rho : G_K \rightarrow \mathrm{GL}_2(A)$  be a lift of  $\bar{\rho}$ . By Hensel's lemma, there are  $a, b \in \mathfrak{m}_A$  such that  $\rho(\varphi)$  has characteristic polynomial  $(X - (\alpha + a))(X - (\beta + b))$ . Show that there are  $x, y \in \mathfrak{m}_A$  such that

$$\rho(\varphi) \begin{pmatrix} 1 \\ x \end{pmatrix} = (\alpha + a) \begin{pmatrix} 1 \\ x \end{pmatrix}$$

and

$$\rho(\varphi) \begin{pmatrix} y \\ 1 \end{pmatrix} = (\beta + b) \begin{pmatrix} y \\ 1 \end{pmatrix}$$

- (2) Since  $\bar{\rho}$  is unramified,  $\bar{\rho}(\sigma) = 1$ , so we may write

$$\begin{pmatrix} 1 & y \\ x & 1 \end{pmatrix}^{-1} \rho(\sigma) \begin{pmatrix} 1 & y \\ x & 1 \end{pmatrix} = \begin{pmatrix} 1 + u & v \\ w & 1 + z \end{pmatrix}$$

with  $u, v, w, z \in \mathfrak{m}_A$ . Use the commutation relation between  $\rho(\varphi)$  and  $\rho(\sigma)$  to show that  $v = w = 0$ .

- (3) Use the fact that  $\chi$  is unramified to show that  $1 + z = (1 + u)^{-1}$ .  
 (4) Show that  $(1 + u)^{\#k} = 1 + u$ , and deduce that  $(1 + u)^{\#k-1} = 1$ .  
 (5) Deduce that  $(1 + u)^{p^m} = 1$ .  
 (6) Complete the proof of the lemma.

**3.35. Taylor's "Ihara avoidance" deformations.** The following deformation rings are crucial to Taylor's arguments in [Tay08] which avoid the use of Ihara's lemma in proving automorphy lifting theorems. When  $n = 2$  these arguments are not logically necessary, but they are crucial to all applications of automorphy lifting theorems when  $n > 2$ .

Continue to let  $K/Q_l$  be a finite extension, and assume that  $\bar{\rho}$  is the trivial 2-dimensional representation, that  $\#k \equiv 1 \pmod{p}$ , that  $\chi$  is unramified, and that  $\bar{\chi}$  is trivial. Again, we see that  $\rho^\square(P_K)$  is trivial, so that  $\rho^\square$  is determined by the two matrices  $\rho^\square(\sigma)$  and  $\rho^\square(\varphi)$ , as in Exercise 3.34. A similar analysis then yields the following facts. (For the proof of the analogous results in the  $n$ -dimensional case, see Section 3 of [Tay08].)

- 3.36. Definition.** (1) Let  $\mathcal{P}_{\mathrm{ur}}$  be the minimal ideal of  $R_{\bar{\rho}, \chi}^\square$  modulo which  $\rho^\square(\sigma) = 1_2$ .  
 (2) For any root of unity  $\zeta$ , we let  $\mathcal{P}_\zeta$  be the minimal ideal of  $R_{\bar{\rho}, \chi}^\square$  modulo which  $\rho^\square(\sigma)$  has characteristic polynomial  $(X - \zeta)(X - \zeta^{-1})$ .  
 (3) Let  $\mathcal{P}_{\mathrm{m}}$  be the minimal ideal of  $R_{\bar{\rho}, \chi}^\square$  modulo which  $\rho^\square(\sigma)$  has characteristic polynomial  $(X - 1)^2$ , and  $\#k(\mathrm{tr} \rho^\square(\varphi))^2 = (1 + \#k)^2 \det \rho^\square(\varphi)$ .

[The motivation for the definition of  $\mathcal{P}_m$  is that we are attempting to describe the unipotent liftings, and if you assume that  $\rho^\square(\sigma) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ , this is the relation forced on  $\rho^\square(\varphi)$ .]

**3.37. Proposition.** *The minimal primes of  $R_{\bar{\rho}, \chi}^\square$  are precisely  $\sqrt{\mathcal{P}_{\text{ur}}}$ ,  $\sqrt{\mathcal{P}_m}$ , and the  $\sqrt{\mathcal{P}_\zeta}$  for  $\zeta \neq 1$ . We have  $\sqrt{\mathcal{P}_1} = \sqrt{\mathcal{P}_{\text{ur}}} \cap \sqrt{\mathcal{P}_m}$ .*

Write  $R_{\bar{\rho}, \chi, 1}^\square$ ,  $R_{\bar{\rho}, \chi, \zeta}^\square$ ,  $R_{\bar{\rho}, \chi, \text{ur}}^\square$ ,  $R_{\bar{\rho}, \chi, m}^\square$  for the corresponding quotients of  $R_{\bar{\rho}, \chi}^\square$ .

**3.38. Theorem.** *We have  $R_{\bar{\rho}, \chi, 1}^\square / \lambda = R_{\bar{\rho}, \chi, \zeta}^\square / \lambda$ . Furthermore,*

- (1)  $R_{\bar{\rho}, \chi, \zeta}^\square[1/p]$  is formally smooth and geometrically irreducible of dimension 3.
- (2)  $R_{\bar{\rho}, \chi, \text{ur}}^\square$  is formally smooth over  $\mathcal{O}$  (and thus geometrically irreducible) of relative dimension 3.
- (3)  $R_{\bar{\rho}, \chi, m}^\square[1/p]$  is formally smooth and geometrically irreducible of dimension 3.
- (4)  $\text{Spec } R_{\bar{\rho}, \chi, 1}^\square = \text{Spec } R_{\bar{\rho}, \chi, \text{ur}}^\square \cup \text{Spec } R_{\bar{\rho}, \chi, m}^\square$  and  $\text{Spec } R_{\bar{\rho}, \chi, 1}^\square / \lambda = \text{Spec } R_{\bar{\rho}, \chi, \text{ur}}^\square / \lambda \cup \text{Spec } R_{\bar{\rho}, \chi, m}^\square / \lambda$  are both a union of two irreducible components, and have relative dimension 3.

*Proof.* See Proposition 3.1 of [Tay08] for an  $n$ -dimensional version of this result. In the 2-dimensional case it can be proved by explicitly computing equations for the lifting rings; see [Sho13].  $\square$

#### 4. MODULAR AND AUTOMORPHIC FORMS, AND THE LANGLANDS CORRESPONDENCE

**4.1. The local Langlands correspondence (and the Jacquet–Langlands correspondence).** Weil–Deligne representations are the objects on the “Galois” side of the local Langlands correspondence. We now describe the objects on the “automorphic” side. These will be representations  $(\pi, V)$  of  $\text{GL}_n(K)$  on (usually infinite-dimensional)  $\mathbb{C}$ -vector spaces.

**4.2. Definition.** We say that  $(\pi, V)$  is *smooth* if for any vector  $v \in V$ , the stabiliser of  $v$  in  $\text{GL}_n(K)$  is open. We say that  $(\pi, V)$  is *admissible* if it is smooth, and for any open subgroup  $U \subset V$ ,  $V^U$  is finite-dimensional.

For example, a smooth one-dimensional representation of  $K^\times$  is the same thing as a continuous character.

- 4.3. Fact.**
- (1) If  $\pi$  is smooth and irreducible then it is admissible.
  - (2) Schur’s lemma is true, and in particular if  $\pi$  is smooth, admissible and irreducible then it has a central character  $\chi_\pi : K^\times \rightarrow \mathbb{C}^\times$ .

In general these representations are classified in terms of the (super)cuspidal representations. We won’t need the details of this classification, and accordingly we won’t define the cuspidal representations.



Let  $B$  be the subgroup of  $\mathrm{GL}_2(K)$  consisting of upper-triangular matrices. Define  $\delta : B \rightarrow K^\times$  by

$$\delta \left( \begin{pmatrix} a & * \\ 0 & d \end{pmatrix} \right) = ad^{-1}.$$

Given two characters  $\chi_1, \chi_2 : K^\times \rightarrow \mathbb{C}^\times$ , we may view  $\chi_1 \otimes \chi_2$  as a representation of  $B$  by

$$\chi_1 \otimes \chi_2 : \begin{pmatrix} a & * \\ 0 & d \end{pmatrix} \mapsto \chi_1(a)\chi_2(d).$$

Then we define a representation  $\chi_1 \times \chi_2$  of  $\mathrm{GL}_2(K)$  by

$$\begin{aligned} \chi_1 \times \chi_2 &= \mathrm{n}\text{-Ind}_B^{\mathrm{GL}_2(K)}(\chi_1 \otimes \chi_2) \\ &:= \{ \varphi : \mathrm{GL}_2(K) \rightarrow \mathbb{C} \mid \varphi(hg) = (\chi_1 \otimes \chi_2)(h) |\delta(h)|_K^{1/2} \varphi(g) \text{ for all } h \in B, g \in \mathrm{GL}_2(K) \} \end{aligned}$$

where  $\mathrm{GL}_2(K)$  acts by  $(g\varphi)(g') = \varphi(g'g)$ , and we only allow smooth  $\varphi$ , i.e. functions for which there is an open subgroup  $U$  of  $\mathrm{GL}_2(K)$  such that  $\varphi(gu) = \varphi(g)$  for all  $g \in \mathrm{GL}_2(K)$ ,  $u \in U$ .

The representation  $\chi_1 \times \chi_2$  has length at most 2, but is not always irreducible. It is always the case that  $\chi_1 \times \chi_2$  and  $\chi_2 \times \chi_1$  have the same Jordan-Hölder factors. If  $\chi_1 \times \chi_2$  is irreducible then we say that it is a *principal series* representation.

4.4. *Fact.* (1)  $\chi_1 \times \chi_2$  is irreducible unless  $\chi_1/\chi_2 = |\cdot|^\pm$ .

(2)  $\chi \times \chi|\cdot|$  has a one-dimensional irreducible subrepresentation, and the corresponding quotient is irreducible. We denote this quotient by  $\mathrm{Sp}_2(\chi)$ .

We will let  $\chi_1 \boxplus \chi_2$  denote  $\chi_1 \times \chi_2$  unless  $\chi_1/\chi_2 = |\cdot|^\pm$ , and we let  $\chi \boxplus \chi|\cdot| = \chi|\cdot| \boxplus \chi = (\chi|\cdot|^{1/2}) \circ \det$ . (While this notation may seem excessive, we remark that a similar construction is possible for  $n$ -dimensional representations, which is where the notation comes from.) These representations, and the  $\mathrm{Sp}_2(\chi)$ , are all the non-cuspidal irreducible admissible representations of  $\mathrm{GL}_2(K)$ . We say that an irreducible smooth representation  $\pi$  of  $\mathrm{GL}_2(K)$  is *discrete series* if it is of the form  $\mathrm{Sp}_2(\chi)$  or is cuspidal.

The local Langlands correspondence provides a unique family of bijections  $\mathrm{rec}_K$  from the set of irreducible smooth representations of  $\mathrm{GL}_n(K)$  to the set of  $n$ -dimensional Frobenius semisimple Weil-Deligne representations of  $W_K$  over  $\mathbb{C}$ , satisfying a list of properties. In order to be uniquely determined, one needs to formulate the correspondence for all  $n$  at once, and the properties are expressed in terms of  $L$ - and  $\varepsilon$ -factors, neither of which we have defined. Accordingly, we will not make a complete statement of the local Langlands correspondence, but will rather state the properties of the correspondence that we will need to use. It is also possible to define the correspondence in global terms, as we will see later, and indeed at present the only proof of the correspondence is global.

4.5. *Fact.* We now list some properties of  $\mathrm{rec}_K$  for  $n = 1, 2$ .

(1) If  $n = 1$  then  $\mathrm{rec}_K(\pi) = \pi \circ \mathrm{Art}_K^{-1}$ .

(2) If  $\chi$  is a smooth character,  $\mathrm{rec}_K(\pi \otimes (\chi \circ \det)) = \mathrm{rec}_K(\pi) \otimes \mathrm{rec}_K(\chi)$ .

- (3)  $\text{rec}_K(\text{Sp}_2(\chi)) = \text{Sp}_2(\text{rec}_K(\chi))$ .
- (4)  $\text{rec}_K(\chi_1 \boxplus \chi_2) = \text{rec}_K(\chi_1) \oplus \text{rec}_K(\chi_2)$ .
- (5)  $\text{rec}_K(\pi)$  is unramified (i.e.  $N = 0$  and the restriction to  $I_K$  is trivial) if and only if  $\pi = \chi_1 \boxplus \chi_2$  with  $\chi_1, \chi_2$  both unramified characters (i.e. trivial on  $\mathcal{O}_K^\times$ ). These conditions are equivalent to  $\pi^{\text{GL}_2(\mathcal{O}_K)} \neq 0$ , in which case it is one-dimensional.
- (6)  $\pi$  is discrete series if and only if  $\text{rec}_K(\pi)$  is indecomposable, and cuspidal if and only if  $\text{rec}_K(\pi)$  is irreducible.

**4.6. Hecke operators.** Let  $\varphi$  be a compactly supported  $\mathbb{C}$ -valued function on  $\text{GL}_2(\mathcal{O}_K) \backslash \text{GL}_2(K) / \text{GL}_2(\mathcal{O}_K)$ . Concretely, these are functions which vanish outside of a finite number of double cosets  $\text{GL}_2(\mathcal{O}_K)g \text{GL}_2(\mathcal{O}_K)$ . The set of such functions is in fact a ring, with the multiplication being given by convolution. To be precise, we fix  $\mu$  the (left and right) Haar measure on  $\text{GL}_2(K)$  such that  $\mu(\text{GL}_2(\mathcal{O}_K)) = 1$ , and we define

$$(\varphi_1 * \varphi_2)(x) = \int_{\text{GL}_2(K)} \varphi_1(g) \varphi_2(g^{-1}x) d\mu_g.$$

Of course, this integral is really just a finite sum. One can check without too much difficulty that the ring  $\mathcal{H}$  of these Hecke operators is just  $\mathbb{C}[T, S^{\pm 1}]$ , where  $T$  is the characteristic function of

$$\text{GL}_2(\mathcal{O}_K) \begin{pmatrix} \varpi_K & 0 \\ 0 & 1 \end{pmatrix} \text{GL}_2(\mathcal{O}_K)$$

and  $S$  is the characteristic function of

$$\text{GL}_2(\mathcal{O}_K) \begin{pmatrix} \varpi_K & 0 \\ 0 & \varpi_K \end{pmatrix} \text{GL}_2(\mathcal{O}_K).$$

The algebra  $\mathcal{H}$  acts on an irreducible admissible  $\text{GL}_2(K)$ -representation  $\pi$ . Given  $\varphi \in \mathcal{H}$ , we obtain a linear map  $\pi(\varphi) : \pi \rightarrow \pi^{\text{GL}_2(\mathcal{O}_K)}$ , by

$$\pi(\varphi)(v) = \int_{\text{GL}_2(K)} \varphi(g) \pi(g)v d\mu_g.$$

In particular, if  $\pi$  is unramified then  $\pi(\varphi)$  acts via a scalar on the one-dimensional  $\mathbb{C}$ -vector space  $\pi^{\text{GL}_2(\mathcal{O}_K)}$ . We will now compute this scalar explicitly.

**4.7. Exercise.** (1) Show that we have decompositions

$$\text{GL}_2(\mathcal{O}_K) \begin{pmatrix} \varpi_K & 0 \\ 0 & \varpi_K \end{pmatrix} \text{GL}_2(\mathcal{O}_K) = \begin{pmatrix} \varpi_K & 0 \\ 0 & \varpi_K \end{pmatrix} \text{GL}_2(\mathcal{O}_K),$$

and

$$\text{GL}_2(\mathcal{O}_K) \begin{pmatrix} \varpi_K & 0 \\ 0 & 1 \end{pmatrix} \text{GL}_2(\mathcal{O}_K) = \left( \prod_{\alpha \in \mathcal{O}_K \atop (\text{mod } \varpi_K)} \begin{pmatrix} \varpi_K & \alpha \\ 0 & 1 \end{pmatrix} \text{GL}_2(\mathcal{O}_K) \right) \prod \begin{pmatrix} 1 & 0 \\ 0 & \varpi_K \end{pmatrix} \text{GL}_2(\mathcal{O}_K).$$

- (2) Suppose that  $\pi = (\chi | \cdot |^{1/2}) \circ \det$  with  $\chi$  unramified. Show that  $\pi^{\text{GL}_2(\mathcal{O}_K)} = \pi$ , and that  $S$  acts via  $\chi(\varpi_K)^2 (\#k)^{-1}$ , and that  $T$  acts via  $(\#k^{1/2} + \#k^{-1/2})\chi(\varpi_K)$ .

- (3) Suppose that  $\chi_1, \chi_2$  are unramified characters and that  $\chi_1 \neq \chi_2 | \cdot |_{\mathbb{K}}^{\pm 1}$ . Let  $\pi = \chi_1 \boxplus \chi_2$ . Using the Iwasawa decomposition  $\mathrm{GL}_2(\mathbb{K}) = B(\mathbb{K}) \mathrm{GL}_2(\mathcal{O}_{\mathbb{K}})$ , check that  $\pi^{\mathrm{GL}_2(\mathcal{O}_{\mathbb{K}})}$  is one-dimensional, and is spanned by a function  $\varphi_0$  with  $\varphi_0 \left( \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \right) = \chi_1(a) \chi_2(d) |a/d|^{1/2}$ . Show that  $S$  acts on  $\pi^{\mathrm{GL}_2(\mathcal{O}_{\mathbb{K}})}$  via  $(\chi_1 \chi_2)(\varpi_{\mathbb{K}})$ , and that  $T$  acts via  $\#k^{1/2}(\chi_1(\varpi_{\mathbb{K}}) + \chi_2(\varpi_{\mathbb{K}}))$ .

**4.8. Modular forms and automorphic forms on quaternion algebras.** Let  $F$  be a totally real field, and let  $D/F$  be a quaternion algebra with centre  $F$ , i.e. central simple  $F$ -algebra of dimension 4. Letting  $S(D)$  be the set of places  $v$  of  $F$  at which  $D$  is ramified, i.e. for which  $D \otimes_F F_v$  is a division algebra (equivalently, is not isomorphic to  $M_2(F_v)$ ), it is known that  $S(D)$  classifies  $D$  up to isomorphism, and that  $S(D)$  can be any finite set of places of  $D$  of even cardinality (so for example  $S(D)$  is empty if and only if  $D = M_2(F)$ ). We will now define some spaces of automorphic forms on  $D^\times$ .

For each  $v|\infty$  fix  $k_v \geq 2$  and  $\eta_v \in \mathbb{Z}$  such that  $k_v + 2\eta_v - 1 = w$  is independent of  $v$ . These will be the weights of our modular forms. Let  $G_D$  be the algebraic group over  $\mathbb{Q}$  such that for any  $\mathbb{Q}$ -algebra  $R$ ,  $G_D(R) = (D \otimes_{\mathbb{Q}} R)^\times$ . For each place  $v|\infty$  of  $F$ , we define a subgroup  $U_v$  of  $(D \otimes_F F_v)^\times$  as follows: if  $v \in S(D)$  we let  $U_v = (D \otimes_F F_v)^\times \cong \mathbb{H}^\times$ , and if  $v \notin S(D)$ , so that  $(D \otimes_F F_v)^\times \cong \mathrm{GL}_2(\mathbb{R})$ , we take  $U_v = \mathbb{R}^\times \mathrm{SO}(2)$ . If  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{R})$  and  $z \in \mathbb{C} - \mathbb{R}$ , we let  $j(\gamma, z) = cz + d$ . One checks easily that  $j(\gamma\delta, z) = j(\gamma, \delta z)j(\delta, z)$ .

We now define a representation  $(\tau_v, W_v)$  of  $U_v$  over  $\mathbb{C}$  for each  $v|\infty$ . If  $v \in S(D)$ , we have  $U_v \hookrightarrow \mathrm{GL}_2(\overline{F}_v) \cong \mathrm{GL}_2(\mathbb{C})$  which acts on  $\mathbb{C}^2$ , and we let  $(\tau_v, W_v)$  be the representation

$$(\mathrm{Sym}^{k_v-2} \mathbb{C}^2) \otimes (\wedge^2 \mathbb{C}^2)^{\eta_v}.$$

If  $v \notin S(D)$ , then we have  $U_v \cong \mathbb{R}^\times \mathrm{SO}(2)$ , and we take  $W_v = \mathbb{C}$ , with

$$\tau_v(\gamma) = j(\gamma, i)^{k_v} (\det \gamma)^{\eta_v-1}.$$

We write  $U_\infty = \prod_{v|\infty} U_v$ ,  $W_\infty = \otimes_{v|\infty} W_v$ ,  $\tau_\infty = \otimes_{v|\infty} \tau_v$ . Let  $\mathbb{A} = \mathbb{A}_{\mathbb{Q}}$  be the adèles of  $\mathbb{Q}$ , and let  $\mathbb{A}^\infty$  be the finite adèles. We then define  $S_{D,k,\eta}$  to be the space of functions  $\varphi : G_D(\mathbb{Q}) \backslash G_D(\mathbb{A}) \rightarrow W_\infty$  which satisfy

- (1)  $\varphi(gu_\infty) = \tau_\infty(u_\infty)^{-1} \varphi(g)$  for all  $u_\infty \in U_\infty$  and  $g \in G_D(\mathbb{A})$ .
- (2) There is a nonempty open subset  $U^\infty \subset G_D(\mathbb{A}^\infty)$  such that  $\varphi(gu) = \varphi(g)$  for all  $u \in U^\infty, g \in G_D(\mathbb{A})$ .
- (3) Let  $S_\infty$  denote the infinite places of  $F$ . If  $g \in G_D(\mathbb{A}^\infty)$  then the function

$$(\mathbb{C} - \mathbb{R})^{S_\infty - S(D)} \rightarrow W_\infty$$

defined by

$$h_\infty(i, \dots, i) \mapsto \tau_\infty(h_\infty) \phi(gh_\infty)$$

is holomorphic. [Note that this function is well-defined by the first condition, as  $U_\infty$  is the stabiliser of  $(i, \dots, i)$ .]

(4) If  $S(D) = \emptyset$  then for all  $g \in G_D(\mathbb{A}) = \mathrm{GL}_2(\mathbb{A}_F)$ , we have

$$\int_{F \backslash \mathbb{A}_F} \varphi \left( \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g \right) dx = 0.$$

If in addition we have  $F = \mathbb{Q}$ , then we furthermore demand that for all  $g \in G_D(\mathbb{A}^\infty)$ ,  $h_\infty \in \mathrm{GL}_2(\mathbb{R})^+$  the function  $\varphi(g h_\infty) |\mathrm{Im}(h_\infty i)|^{k/2}$  is bounded on  $\mathbb{C} - \mathbb{R}$ .

There is a natural action of  $G_D(\mathbb{A}^\infty)$  on  $S_{D,k,\eta}$  by right-translation, i.e.  $(g\varphi)(x) := \varphi(xg)$ .

4.9. *Exercise.* While this definition may at first sight appear rather mysterious, it is just a generalisation of the familiar spaces of cuspidal modular forms. For example, take  $F = \mathbb{Q}$ ,  $S(D) = \emptyset$ ,  $k_\infty = k$ , and  $\eta_\infty = 0$ . Define

$$U_1(N) = \{g \in \mathrm{GL}_2(\widehat{\mathbb{Z}}) \mid g \equiv \begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix} \pmod{N}\}.$$

- (1) Let  $\mathrm{GL}_2(\mathbb{Q})^+$  be the subgroup of  $\mathrm{GL}_2(\mathbb{Q})$  consisting of matrices with positive determinant. Show that the intersection of  $\mathrm{GL}_2(\mathbb{Q})^+$  and  $U_1(N)$  inside  $\mathrm{GL}_2(\mathbb{A}^\infty)$  is  $\Gamma_1(N)$ , the matrices in  $\mathrm{SL}_2(\mathbb{Z})$  congruent to  $\begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \pmod{N}$ . [Hint: what is  $\widehat{\mathbb{Z}}^\times \cap \mathbb{Q}^\times$ ?]
- (2) Use the facts that  $\mathrm{GL}_2(\mathbb{A}) = \mathrm{GL}_2(\mathbb{Q})U_1(N)\mathrm{GL}_2(\mathbb{R})^+$  [which follows from strong approximation for  $\mathrm{SL}_2$  and the fact that  $\det U_1(N) = \widehat{\mathbb{Z}}^\times$ ] and that  $\mathbb{A}^\times = \mathbb{Q}^\times \widehat{\mathbb{Z}}^\times \mathbb{R}_{>0}^\times$  to show that  $S_{D,k,0}^{U_1(N)}$  can naturally be identified with a space of functions

$$\varphi : \Gamma_1(N) \backslash \mathrm{GL}_2(\mathbb{R})^+ \rightarrow \mathbb{C}$$

satisfying

$$\varphi(g u_\infty) = j(u_\infty, i)^{-k} \varphi(g)$$

for all  $g \in \mathrm{GL}_2(\mathbb{R})^+$ ,  $u_\infty \in \mathbb{R}_{>0}^\times \mathrm{SO}(2)$ .

- (3) Show that the stabiliser of  $i$  in  $\mathrm{GL}_2(\mathbb{R})^+$  is  $\mathbb{R}_{>0}^\times \mathrm{SO}(2)$ . Hence deduce a natural isomorphism between  $S_{D,k,0}^{U_1(N)}$  and  $S_k(\Gamma_1(N))$ , which takes a function  $\varphi$  as above to the function  $(gi \mapsto j(g, i)^k \varphi(g))$ ,  $g \in \mathrm{GL}_2(\mathbb{R})^+$ .

The case that  $S_\infty \subset S(D)$  is particularly simple; then if  $U \subset G_D(\mathbb{A}^\infty)$  is an open subgroup, then  $S_{D,2,0}^U$  is just the set of  $\mathbb{C}$ -valued functions on

$$G_D(\mathbb{Q}) \backslash G_D(\mathbb{A}) / G_D(\mathbb{R})U,$$

which is a finite set. When proving modularity lifting theorems, we will be able to reduce to the case that  $S_\infty \subset S(D)$ ; when this condition holds, we say that  $D$  is a *definite* quaternion algebra.

We will now examine the action of Hecke operators on these spaces. Choose an order  $\mathcal{O}_D \subset D$  (that is, a  $\mathbb{Z}$ -subalgebra of  $D$  which is finitely generated as a  $\mathbb{Z}$ -module and for which  $\mathcal{O}_D \otimes_{\mathbb{Z}} \mathbb{Q} \xrightarrow{\sim} D$ ). For example, if  $D = M_2(F)$ , one may take  $\mathcal{O}_D = M_2(\mathcal{O}_F)$ .

For all but finite many finite places  $v$  of  $F$  we can choose an isomorphism  $D_v \cong M_2(F_v)$  such that this isomorphism induces an isomorphism  $\mathcal{O}_D \otimes_{\mathcal{O}_F} \mathcal{O}_{F_v} \xrightarrow{\sim} M_2(\mathcal{O}_{F_v})$ . Then  $G_D(\mathbb{A}^\infty)$  is the subset of elements  $g = (g_v) \in \prod_{v \neq \infty} G_D(F_v)$  such that  $g_v \in \mathrm{GL}_2(\mathcal{O}_{F_v})$  for almost all  $v$ .

We now wish to describe certain irreducible representations of  $G_D(\mathbb{A}^\infty)$  in terms of irreducible representations of the  $\mathrm{GL}_2(F_v)$ . More generally, we have the following construction. Let  $I$  be an indexing set and let  $V_i$  be a  $\mathbb{C}$ -vector space. Suppose that we are given  $0 \neq e_i \in V_i$  for almost all  $i$  (that is, all but finitely many  $i$ ). Then we define the *restricted tensor product*

$$\otimes'_{\{e_i\}} V_i := \varinjlim_{J \subset I} \otimes_{i \in J} V_i,$$

where the colimit is over the finite subsets  $J \subseteq I$  containing all the places for which  $e_i$  is not defined, and where the transition maps for the colimit are given by “tensoring with the  $e_i$ ”. It can be checked that  $\otimes'_{\{e_i\}} V_i \cong \otimes'_{\{f_i\}} V_i$  if for almost all  $i$ ,  $e_i$  and  $f_i$  span the same line.

**4.10. Definition.** We call a representation  $(\pi, V)$  of  $G_D(\mathbb{A}^\infty)$  *admissible* if

- (1) for any  $x \in V$ , the stabiliser of  $x$  is open, and
- (2) for any  $U \subset G_D(\mathbb{A}^\infty)$  an open subgroup,  $\dim_{\mathbb{C}} V^U < \infty$ .

**4.11. Fact.** If  $\pi_v$  is an irreducible smooth (so admissible) representation of  $(D \otimes_F F_v)^\times$  with  $\pi_v^{\mathrm{GL}_2(\mathcal{O}_{F_v})} \neq 0$  for almost all  $v$ , then  $\otimes' \pi_v := \otimes'_{\left\{ \pi_v^{\mathrm{GL}_2(\mathcal{O}_{F_v})} \right\}} \pi_v$  is an irreducible admissible

smooth representation of  $G_D(\mathbb{A}^\infty)$ , and any irreducible admissible smooth representation of  $G_D(\mathbb{A}^\infty)$  arises in this way for unique  $\pi_v$ .

We have a global Hecke algebra, which decomposes as a restricted product of the local Hecke algebras in the following way. For each finite place  $v$  of  $F$  we choose  $U_v \subset G_D(F_v)$  a compact open subgroup, such that  $U_v = \mathrm{GL}_2(\mathcal{O}_{F_v})$  for almost all  $v$ . Let  $\mu_v$  be a Haar measure on  $G_D(F_v)$ , chosen such that for almost all  $v$  we have  $\mu_v(\mathrm{GL}_2(\mathcal{O}_{F_v})) = 1$ . Then there is a unique Haar measure  $\mu$  on  $G_D(\mathbb{A}^\infty)$  such that for any  $U_v$  as above, if we set  $U = \prod_v U_v \subset G_D(\mathbb{A}^\infty)$ , then  $\mu(U) = \prod_v \mu_v(U_v)$ . Then there is a decomposition

$$\mathcal{C}_c(U \backslash G_D(\mathbb{A}^\infty) / U) \mu \cong \otimes'_{\{1_{U_v \mu_v}\}} \mathcal{C}_c(U_v \backslash G_D(F_v) / U_v) \mu_v,$$

and the actions of these Hecke algebras are compatible with the decomposition  $\pi = \otimes' \pi_v$ .

**4.12. Fact.**  $S_{D,k,\eta}$  is a semisimple admissible representation of  $G_D(\mathbb{A}^\infty)$ .

**4.13. Definition.** The irreducible constituents of  $S_{D,k,\eta}$  are called the *cuspidal automorphic representations* of  $G_D(\mathbb{A}^\infty)$  of weight  $(k, \eta)$ .

4.14. *Remark.* Note that these automorphic representations do not include Maass forms or weight one modular forms; they are the class of *regular algebraic* or *cohomological* cuspidal automorphic representations.

4.15. *Fact.* (Strong multiplicity one (and multiplicity one) for  $GL_2$ ) Suppose that  $S(D) = \emptyset$ . Then every irreducible constituent of  $S_{D,k,\eta}$  has multiplicity one. In fact if  $\pi$  (respectively  $\pi'$ ) is a cuspidal automorphic representation of weight  $(k, \eta)$  (respectively  $(k', \eta')$ ) such that  $\pi_v \cong \pi'_v$  for almost all  $v$  then  $k = k'$ ,  $\eta = \eta'$ , and  $\pi = \pi'$ .

4.16. *Fact.* (The theory of newforms) Suppose that  $S(D) = \emptyset$ . If  $\mathfrak{n}$  is an ideal of  $\mathcal{O}_F$ , write

$$U_1(\mathfrak{n}) = \{g \in GL_2(\hat{\mathcal{O}}_F) \mid g \equiv \begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix} \pmod{\mathfrak{n}}\}.$$

If  $\pi$  is a cuspidal automorphic representation of  $G_D(\mathbb{A}^\infty)$  then there is a unique ideal  $\mathfrak{n}$  such that  $\pi^{U_1(\mathfrak{n})}$  is one-dimensional, and  $\pi^{U_1(\mathfrak{m})} \neq 0$  if and only if  $\mathfrak{n} \mid \mathfrak{m}$ . We call  $\mathfrak{n}$  the *conductor* (or sometimes the *level*) of  $\pi$ .

Analogous to the theory of admissible representations of  $GL_2(K)$ ,  $K/\mathbb{Q}_p$  finite that we sketched above, there is a theory of admissible representations of  $D^\times$ ,  $D$  a nonsplit quaternion algebra over  $K$ . Since  $D^\times/K^\times$  is compact, any irreducible smooth representation of  $D^\times$  is finite-dimensional. There is a bijection  $JL$ , the *local Jacquet–Langlands correspondence*, from the irreducible smooth representations of  $D^\times$  to the discrete series representations of  $GL_2(K)$ , determined by a character identity.

4.17. *Fact* (The global Jacquet–Langlands correspondence). We have the following facts about  $G_D(\mathbb{A}^\infty)$ .

- (1) The only finite-dimensional cuspidal automorphic representations of  $G_D(\mathbb{A}^\infty)$  are 1-dimensional representations which factor through the reduced determinant; these only exist if  $G_D \neq GL_2$ .
- (2) There is a bijection  $JL$  from the infinite-dimensional cuspidal automorphic representations of  $G_D(\mathbb{A}^\infty)$  of weight  $(k, \eta)$  to the cuspidal automorphic representations of  $GL_2(\mathbb{A}_F^\infty)$  of weight  $(k, \eta)$  which are discrete series for all finite places  $v \in S(D)$ . Furthermore if  $v \notin S(D)$  then  $JL(\pi)_v = \pi_v$ , and if  $v \in S(D)$  then  $JL(\pi)_v = JL(\pi_v)$ .

4.18. *Remark.* We will use the global Jacquet–Langlands correspondence together with base change (see below) to reduce ourselves to considering the case that  $S(D) = S_\infty$  when proving automorphy lifting theorems.

#### 4.19. Galois representations associated to automorphic representations.

4.20. *Fact* (The existence of Galois representations associated to regular algebraic cuspidal automorphic representations). Let  $\pi$  be a regular algebraic cuspidal automorphic representation

of  $\mathrm{GL}_2(\mathbb{A}_F^\infty)$  of weight  $(k, \eta)$ . Then there is a CM field  $L_\pi$  and for each finite place  $\lambda$  of  $L_\pi$  a continuous irreducible Galois representation

$$r_\lambda(\pi) : G_F \rightarrow \mathrm{GL}_2(\overline{L}_{\pi, \lambda})$$

such that

- (1) if  $\pi_v$  is unramified and  $v$  does not divide the residue characteristic of  $\lambda$ , then  $r_\lambda(\pi)|_{G_{F_v}}$  is unramified, and the characteristic polynomial of  $\mathrm{Frob}_v$  is  $X^2 - t_v X + (\#k_v)s_v$ , where  $t_v$  and  $s_v$  are the eigenvalues of  $T_v$  and  $S_v$  respectively on  $\pi_v^{\mathrm{GL}_2(\mathcal{O}_{F_v})}$ . [Note that by the Chebotarev density theorem, this already characterises  $r_\lambda(\pi)$  up to isomorphism.]
- (2) More generally,  $\mathrm{WD}(r_\lambda(\pi)|_{G_{F_v}})^{F\text{-ss}} \cong \mathrm{rec}_{F_v}(\pi_v \otimes |\det|^{-1/2})$ .
- (3) If  $v$  divides the residue characteristic of  $\lambda$  then  $r_\lambda(\pi)|_{G_{F_v}}$  is de Rham with  $\tau$ -Hodge-Tate weights  $\eta_\tau, \eta_\tau + k_\tau - 1$ , where  $\tau : F \hookrightarrow \overline{L}_\pi \subset \mathbb{C}$  is an embedding lying over  $v$ . If  $\pi_v$  is unramified then  $r_\lambda(\pi)|_{G_{F_v}}$  is crystalline.
- (4) If  $c_v$  is a complex conjugation, then  $\det r_\lambda(\pi)(c_v) = -1$ .

4.21. *Remark.* Using the Jacquet–Langlands correspondence, we get Galois representations for the infinite-dimensional cuspidal automorphic representations of  $G_D(\mathbb{A}^\infty)$  for any  $D$ . In fact, the proof actually uses the Jacquet–Langlands correspondence; in most cases, you can transfer to a  $D$  for which  $S(D)$  contains all but one infinite place, and the Galois representations are then realised in the étale cohomology of the associated Shimura curve. The remaining Galois representations are constructed from these ones via congruences.

4.22. *Fact* (Cyclic base change). Let  $E/F$  be a cyclic extension of totally real fields of prime degree. Let  $\mathrm{Gal}(E/F) = \langle \sigma \rangle$  and let  $\mathrm{Gal}(E/F)^\vee = \langle \delta_{E/F} \rangle$  (here  $\mathrm{Gal}(E/F)^\vee$  is the dual abelian group of  $\mathrm{Gal}(E/F)$ ). Let  $\pi$  be a cuspidal automorphic representation of  $\mathrm{GL}_2(\mathbb{A}_F^\infty)$  of weight  $(k, \eta)$ . Then there is a cuspidal automorphic representation  $\mathrm{BC}_{E/F}(\pi)$  of  $\mathrm{GL}_2(\mathbb{A}_E^\infty)$  of weight  $(\mathrm{BC}_{E/F}(k), \mathrm{BC}_{E/F}(\eta))$  such that

- (1) for all finite places  $v$  of  $E$ ,  $\mathrm{rec}_{E_v}(\mathrm{BC}_{E/F}(\pi)_v) = (\mathrm{rec}_{F_v|F}(\pi_{v|F}))|_{W_{E_v}}$ . In particular,  $r_\lambda(\mathrm{BC}_{E/F}(\pi)) \cong r_\lambda(\pi)|_{G_E}$ .
- (2)  $\mathrm{BC}_{E/F}(k)_v = k_{v|F}$ ,  $\mathrm{BC}_{E/F}(\eta)_v = \eta_{v|F}$ .
- (3)  $\mathrm{BC}_{E/F}(\pi) \cong \mathrm{BC}_{E/F}(\pi')$  if and only if  $\pi \cong \pi' \otimes (\delta_{E/F}^i \circ \mathrm{Art}_F \circ \det)$  for some  $i$ .
- (4) A cuspidal automorphic representation  $\pi$  of  $\mathrm{GL}_2(\mathbb{A}_E^\infty)$  is in the image of  $\mathrm{BC}_{E/F}$  if and only if  $\pi \circ \sigma \cong \pi$ .

4.23. **Definition.** We say that  $r : G_F \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}}_p)$  is *modular* (of weight  $(k, \eta)$ ) if it is isomorphic to  $i(r_\lambda(\pi))$  for some cuspidal automorphic representation  $\pi$  (of weight  $(k, \eta)$ ) and some  $i : L_\pi \hookrightarrow \overline{\mathbb{Q}}_p$  lying over  $\lambda$ .

4.24. **Proposition.** *Suppose that  $r : G_F \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}}_p)$  is a continuous representation, and that  $E/F$  is a finite solvable Galois extension of totally real fields. Then  $r|_{G_E}$  is modular if and only if  $r$  is modular.*

4.25. *Exercise.* Prove the above proposition as follows.

- (1) Use induction to reduce to the case that  $E/F$  is cyclic of prime degree.
- (2) Suppose that  $r|_{G_E}$  is modular, say  $r|_{G_E} \cong i(r_\lambda(\pi))$ . Use strong multiplicity one to show that  $\pi \circ \sigma \cong \pi$ . Deduce that there is an automorphic representation  $\pi'$  such that  $\text{BC}_{E/F}(\pi') = \pi$ .
- (3) Use Schur's lemma to deduce that there is a character  $\chi$  of  $G_F$  such that  $r \cong i(r_\lambda(\pi')) \otimes \chi$ . Conclude that  $r$  is modular.

We can make use of this result to make considerable simplifications in our proofs of modularity lifting theorems. It is frequently employed in conjunction with the following fact from class field theory.

4.26. *Fact* (Lemma 2.2 of [Tay03]). Let  $K$  be a number field, and let  $S$  be a finite set of places of  $K$ . For each  $v \in S$ , let  $L_v$  be a finite Galois extension of  $K_v$ . Then there is a finite solvable Galois extension  $M/K$  such that for each place  $w$  of  $M$  above a place  $v \in S$  there is an isomorphism  $L_v \cong M_w$  of  $K_v$ -algebras.

Note that we are allowed to have infinite places in  $S$ , so that if  $K$  is totally real we may choose to make  $L$  totally real by an appropriate choice of the  $L_v$ .

## 5. THE TAYLOR–WILES–KISIN METHOD

Our aim now is to prove the following theorem. Let  $p > 3$  be prime, and let  $L/\mathbb{Q}_p$  be a finite extension with ring of integers  $\mathcal{O}$ , maximal ideal  $\lambda$ , and residue field  $\mathbb{F} = \mathcal{O}/\lambda$ . Let  $F$  be a totally real number field, and assume that  $L$  is sufficiently large that  $L$  contains the images of all embeddings  $F \hookrightarrow \bar{L}$ .

5.1. **Theorem.** *Let  $\rho, \rho_0 : G_F \rightarrow \text{GL}_2(\mathcal{O})$  be two continuous representations, such that  $\bar{\rho} = \rho \pmod{\lambda} = \rho_0 \pmod{\lambda}$ . Assume that  $\rho_0$  is modular, and that  $\rho$  is geometric. Assume further that the following properties hold.*

- (1) *For all  $\sigma : F \hookrightarrow L$ ,  $\text{HT}_\sigma(\rho) = \text{HT}_\sigma(\rho_0)$ , and contains two distinct elements.*
- (2)
  - *For all  $v|p$ ,  $\rho|_{G_{\mathbb{F}_v}}$  and  $\rho_0|_{G_{\mathbb{F}_v}}$  are crystalline.*
  - *$p$  is unramified in  $F$ .*
  - *For all  $\sigma : F \hookrightarrow L$ , the elements of  $\text{HT}_\sigma(\rho)$  differ by at most  $p - 2$ .*
- (3)  *$\text{Im } \bar{\rho} \supseteq \text{SL}_2(\mathbb{F}_p)$ .*

*Then  $\rho$  is modular.*

5.2. **The integral theory of automorphic forms.** In order to prove Theorem 5.1, we will need to study congruences between automorphic forms. In order to do this, it is convenient to work with automorphic forms on  $G_D(\mathbb{A}^\infty)$ , where  $S(D) = S_\infty$ . In order to do this, assume that  $[F : \mathbb{Q}]$  is even. (We will reduce to this case by base change.) Then such a  $D$  exists, and we have  $G_D(\mathbb{A}^\infty) \cong \text{GL}_2(\mathbb{A}_F^\infty)$ , and  $(D \otimes_{\mathbb{Q}} \mathbb{R})^\times / (F \otimes_{\mathbb{Q}} \mathbb{R})^\times$  is compact.



Fix an isomorphism  $\iota : \bar{L} \xrightarrow{\sim} \mathbf{C}$ , and some  $k \in \mathbb{Z}_{\geq 2}^{\text{Hom}(F, \mathbf{C})}$ ,  $\eta \in \mathbb{Z}^{\text{Hom}(F, \mathbf{C})}$  with  $w := k_\tau + 2\eta_\tau - 1$  independent of  $\tau$ . Let  $U = \prod_v U_v \subset \text{GL}_2(\mathbb{A}_F^\infty)$  be a compact open subgroup, and let  $S$  be a finite set of finite places of  $F$ , not containing any of the places lying over  $p$ , with the property that if  $v \notin S$ , then  $U_v = \text{GL}_2(\mathcal{O}_{F_v})$ .

Let  $U_S := \prod_{v \in S} U_v$ , write  $U = U_S U^S$ , let  $\psi : U_S \rightarrow \mathcal{O}^\times$  be a continuous homomorphism (which implies that it has open kernel), and let  $\chi_0 : \mathbb{A}_F^\times / F^\times \rightarrow \mathbf{C}^\times$  be an algebraic grossen-character with the properties that

- $\chi_0$  is unramified outside  $S$ ,
- for each place  $v \mid \infty$ ,  $\chi_0|_{(F_v^\times)^\circ}(x) = x^{1-w}$ , and
- $\chi_0|_{(\prod_{v \in S} F_v^\times) \cap U_S} = \psi^{-1}$ .

As in Theorem 2.41, this gives us a character

$$\chi_{0, \iota} : \mathbb{A}_F^\times / F^\times \overline{(F_\infty^\times)^\circ} \rightarrow \bar{L}^\times,$$

$$x \mapsto \left( \prod_{\tau: \bar{F} \rightarrow L} \tau(x_p)^{1-w} \right) \iota^{-1} \left( \prod_{\tau: \bar{F} \rightarrow \mathbf{C}} \tau(x_\infty) \right)^{w-1} \chi_0(x).$$

Our spaces of ( $p$ -adic) algebraic automorphic forms will be defined in a similar way to the more classical spaces defined in Section 4.8, but with the role of the infinite places being played by the places lying over  $p$ . Accordingly, we define coefficient systems in the following way.

Let  $\Lambda = \Lambda_{k, \eta, \iota} = \otimes_{\tau: F \rightarrow \mathbf{C}} \text{Sym}^{k_\tau - 2}(\mathcal{O}^2) \otimes (\wedge^2 \mathcal{O}^2)^{\otimes \eta_\tau}$ , and let  $\text{GL}_2(\mathcal{O}_{F, p}) := \prod_{v \mid p} \text{GL}_2(\mathcal{O}_{F_v})$  act on  $\Lambda$  via  $\iota^{-1} \tau$  on the  $\tau$ -factor. In particular,  $\Lambda \otimes_{\mathcal{O}, \iota} \mathbf{C} \cong \otimes_{\tau: F \rightarrow \mathbf{C}} \text{Sym}^{k_\tau - 2}(\mathbf{C}^2) \otimes (\wedge^2 \mathbf{C}^2)^{\otimes \eta_\tau}$ , which has an obvious action of  $\text{GL}_2(F_\infty)$ , and the two actions of  $\text{GL}_2(\mathcal{O}_{F, (p)})$  (via its embeddings into  $\text{GL}_2(\mathcal{O}_{F, p})$  and  $\text{GL}_2(F_\infty)$ ) are compatible.

Let  $A$  be finite  $\mathcal{O}$ -module. Then we define  $S(U, A) = S_{k, \eta, \iota, \psi, \chi_0}(U, A)$  to be the spaces of functions

$$\phi : D^\times \backslash \text{GL}_2(\mathbb{A}_F^\infty) \rightarrow \Lambda \otimes_{\mathcal{O}} A$$

such that for all  $g \in \text{GL}_2(\mathbb{A}_F^\infty)$ ,  $u \in U$ ,  $z \in (\mathbb{A}_F^\infty)^\times$ , we have

$$\phi(guz) = \chi_{0, \iota}(z) \psi(u_S)^{-1} u_p^{-1} \phi(g).$$

Since  $D^\times \backslash \text{GL}_2(\mathbb{A}_F^\infty) / U (\mathbb{A}_F^\infty)^\times$  is finite, we see in particular that  $S(U, \mathcal{O})$  is a finite free  $\mathcal{O}$ -module. It has a Hecke action in the obvious way: let  $\tilde{\mathbb{T}} := \mathcal{O}[T_v, S_v : v \nmid p, v \notin S]$ , and let  $T_v, S_v$  act via the usual double coset operators corresponding to  $\begin{pmatrix} \varpi_v & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} \varpi_v & 0 \\ 0 & \varpi_v \end{pmatrix}$ . Let  $\mathbb{T}_U$  be the image of  $\tilde{\mathbb{T}}$  in  $\text{End}_{\mathcal{O}}(S(U, \mathcal{O}))$ , so that  $\mathbb{T}_U$  is a commutative  $\mathcal{O}$ -algebra which acts faithfully on  $S(U, \mathcal{O})$ , and is finite free as an  $\mathcal{O}$ -module.

By definition, we see that

$$S(U, \mathcal{O}) \otimes_{\mathcal{O}, \iota} \mathbf{C} \xrightarrow{\sim} \text{Hom}_{U_S}(\mathbf{C}(\psi^{-1}), S_{k, \eta}^{U^S, \chi_0}),$$

with the map being

$$\phi \mapsto (g \mapsto g_\infty^{-1} \iota(g_p \phi(g^\infty))),$$

and the target of the isomorphism being the elements  $\phi \in S_{k,\eta}$  with  $z\phi = \chi_0(z)\phi$  for all  $z \in (\mathbb{A}_F^\infty)^\times$ ,  $u\phi = \psi(u_S)^{-1}\phi$  for all  $u \in U$ . This isomorphism is compatible with the actions of  $\mathbb{T}$  on each side. The target is isomorphic to

$$\bigoplus_{\pi} \mathrm{Hom}_{U_S}(\mathbb{C}(\psi^{-1}), \pi_S) \otimes \bigotimes'_{v \notin S} \pi_v^{\mathrm{GL}_2(\mathcal{O}_{F_v})},$$

where the sum is over the cuspidal automorphic representations  $\pi$  of  $G_D(\mathbb{A}^\infty)$  of weight  $(k, \eta)$ , which have central character  $\chi_0$  and are unramified outside of  $S$  (so that in particular, for  $v \notin S$ ,  $\pi_v^{\mathrm{GL}_2(\mathcal{O}_{F_v})}$  is a one-dimensional  $\mathbb{C}$ -vector space).

By strong multiplicity one, this means that we have an isomorphism

$$\mathbb{T}_U \otimes_{\mathcal{O}, \iota} \mathbb{C} \cong \prod_{\pi \text{ as above, with } \mathrm{Hom}_{U_S}(\mathbb{C}(\psi^{-1}), \pi_S) \neq (0)} \mathbb{C}$$

sending  $T_v, S_v$  to their eigenvalues on  $\pi_v^{\mathrm{GL}_2(\mathcal{O}_{F_v})}$ . (Note in particular that this shows that  $\mathbb{T}_U$  is reduced.) This shows that there is a bijection between  $\iota$ -linear ring homomorphisms  $\theta : \mathbb{T}_U \rightarrow \mathbb{C}$  and the set of  $\pi$  as above, where  $\pi$  corresponds to the character taking  $T_v, S_v$  to their corresponding eigenvalues.

Each  $\pi$  has a corresponding Galois representation. Taking the product of these representations, we obtain a representation

$$\rho^{\mathrm{mod}} : G_F \rightarrow \prod_{\pi} \mathrm{GL}_2(\bar{L}) = \mathrm{GL}_2(\mathbb{T}_U \otimes_{\mathcal{O}} \bar{L}),$$

which is characterised by the properties that it is unramified outside of  $S \cup \{v|p\}$ , and for any  $v \notin S$ ,  $v \nmid p$ , we have  $\mathrm{tr} \rho^{\mathrm{mod}}(\mathrm{Frob}_v) = T_v$ ,  $\det \rho^{\mathrm{mod}}(\mathrm{Frob}_v) = S_v \# k(v)$ .

Let  $\mathfrak{m}$  be a maximal ideal of  $\mathbb{T}_U$ . Then if  $\mathfrak{p} \subsetneq \mathfrak{m}$  is a minimal prime, then there is an injection  $\theta : \mathbb{T}_U/\mathfrak{p} \hookrightarrow \bar{L}$ , which corresponds to some  $\pi$  as above. (This follows from the going-up and going-down theorems, and the fact that  $\mathbb{T}_U$  is finitely generated and free over  $\mathcal{O}$ .) The semisimple mod  $\mathfrak{p}$  Galois representation corresponding to  $\pi$  can be conjugated to give a representation  $\bar{\rho}_{\mathfrak{m}} : G_F \rightarrow \mathrm{GL}_2(\mathbb{T}_U/\mathfrak{m})$  (because the trace and determinant are valued in  $\mathbb{T}_U/\mathfrak{m}$ , which is a finite field, and thus has trivial Brauer group). This is well defined (up to isomorphism) independently of the choice of  $\mathfrak{p}$  and  $\theta$  (by the Chebotarev density theorem).

Since  $\mathbb{T}_U$  is finite over the complete local ring  $\mathcal{O}$ , it is semilocal, and we can write  $\mathbb{T}_U = \prod_{\mathfrak{m}} \mathbb{T}_{U, \mathfrak{m}}$ . Suppose now that  $\bar{\rho}_{\mathfrak{m}}$  is absolutely irreducible. Then we have the representation

$$\rho_{\mathfrak{m}}^{\mathrm{mod}} : G_F \rightarrow \mathrm{GL}_2(\mathbb{T}_{U, \mathfrak{m}} \otimes_{\mathcal{O}} \bar{L}) = \prod_{\pi} \mathrm{GL}_2(\bar{L}),$$

where the product is over the  $\pi$  as above with  $\bar{\rho}_{\pi, \iota} \cong \bar{\rho}_{\mathfrak{m}}$ . Each representation to  $\mathrm{GL}_2(\bar{L})$  can be conjugated to lie in  $\mathrm{GL}_2(\mathcal{O}_{\bar{L}})$ , and after further conjugation (so that the residual representations are equal to  $\bar{\rho}_{\mathfrak{m}}$ , rather than just conjugate to it), the image of  $\rho_{\mathfrak{m}}^{\mathrm{mod}}$  lies in the subring of  $\prod_{\pi} \mathrm{GL}_2(\mathcal{O}_{\bar{L}})$  consisting of elements whose image modulo the maximal ideal of  $\mathcal{O}_{\bar{L}}$  lie in

$\mathbb{T}_U/\mathfrak{m}$ . We can then apply Lemma 3.7 to see that  $\rho_{\mathfrak{m}}^{\text{mod}}$  can be conjugated to lie in  $\text{GL}_2(\mathbb{T}_{U,\mathfrak{m}})$ . We will write  $\rho_{\mathfrak{m}}^{\text{mod}} : G_F \rightarrow \text{GL}_2(\mathbb{T}_{U,\mathfrak{m}})$  for the resulting representation from now on.

We will sometimes want to consider Hecke operators at places in  $S$ . To this end, let  $T \subseteq S$  satisfy  $\psi|_{U_T} = 1$ , and choose  $g_v \in \text{GL}_2(F_v)$  for each  $v \in T$ . Set  $W_v = [U_v g_v U_v]$ , and define  $\mathbb{T}_U \subseteq \mathbb{T}'_U \subseteq \text{End}_{\mathcal{O}}(S(U, \mathcal{O}))$  by adjoining the  $W_v$  for  $v \in T$ . This is again commutative, and finite and flat over  $\mathcal{O}$ . However, it need not be reduced; indeed, we have

$$\mathbb{T}'_U \otimes_{\mathcal{O},t} \mathbb{C} \cong \oplus_{\pi} \otimes_{v \in T} \{ \text{subalgebra of } \text{End}_{\mathbb{C}}(\pi_v^{U_v}) \text{ generated by } W_v \},$$

so that there is a bijection between  $t$ -linear homomorphisms  $\mathbb{T}'_U \rightarrow \mathbb{C}$  and tuples  $(\pi, \{\alpha_v\}_{v \in T})$ , where  $\alpha_v$  is an eigenvalue of  $W_v$  on  $\pi_v^{U_v}$ . (Note that we will not explicitly use the notation  $\mathbb{T}'_U$  again for a Hecke algebra, but that for example the Hecke algebras  $\mathbb{T}_{U_Q}$  used in the patching argument below, which incorporate Hecke operators at the places in  $Q$ , are an example of this construction.)

We can write

$$\text{GL}_2(\mathbb{A}_F^{\infty}) = \prod_{i \in I} D^{\times} g_i U(\mathbb{A}_F^{\infty})^{\times}$$

for some finite indexing set  $I$ , and so we have an injection  $S(U, A) \hookrightarrow \oplus_{i \in I} (\Lambda \otimes A)$ , by sending  $\phi \mapsto (\phi(g_i))$ . To determine the image, we need to consider when we can have  $g_i = \delta g_i u z$  for  $\delta \in D^{\times}$ ,  $z \in (\mathbb{A}_F^{\infty})^{\times}$ ,  $u \in U$  (because then  $\phi(g_i) = \phi(\delta g_i u z) = \chi_{0,i}(z) \psi(u_S)^{-1} u_p^{-1} \phi(g_i)$ ). We see in this way that we obtain an isomorphism

$$S(U, A) \xrightarrow{\sim} \oplus_{i \in I} (\Lambda \otimes A)^{(U(\mathbb{A}_F^{\infty})^{\times} \cap g_i^{-1} D^{\times} g_i) / F^{\times}}.$$

We need to have some control on these finite groups  $G_i := (U(\mathbb{A}_F^{\infty})^{\times} \cap g_i^{-1} D^{\times} g_i) / F^{\times}$ . (Note that they are finite, because  $D^{\times}$  is discrete in  $G_D(\mathbb{A}^{\infty})$ .) Since we have assumed that  $p > 3$  and  $p$  is unramified in  $F$ , we see that  $[F(\zeta_p) : F] > 2$ . Then we claim that  $G_i$  has order prime to  $p$ . To see this, note that if  $g_i^{-1} \delta g_i$  is in this group, with  $\delta \in D^{\times}$ , then  $\delta^2 / \det \delta \in D^{\times} \cap g_i U g_i^{-1} (\det U)$ , the intersection of a discrete set and a compact set, so  $\delta^2 / \det \delta$  has finite order, i.e. is a root of unity. However any element of  $D$  generates an extension of  $F$  of degree at most 2, so by the assumption that  $[F(\zeta_p) : F] > 2$ , it must be a root of unity of degree prime to  $p$ , and there is some  $p \nmid N$  with  $\delta^{2N} \in F^{\times}$ , so that  $g_i^{-1} \delta g_i$  has order prime to  $p$ , as required.

**5.3. Proposition.** (1) We have  $S(U, \mathcal{O}) \otimes_{\mathcal{O}} A \xrightarrow{\sim} S(U, A)$ .

(2) If  $V$  is an open normal subgroup of  $U$  with  $\#(U/V)$  a power of  $p$ , then  $S(V, \mathcal{O})$  is a free  $\mathcal{O}[U/V(U \cap (\mathbb{A}_F^{\infty})^{\times})]$ -module.

*Proof.* (1) This is immediate from the isomorphism  $S(U, A) \xrightarrow{\sim} \oplus_{i \in I} (\Lambda \otimes A)^{G_i}$ , because the fact that the  $G_i$  have order prime to  $p$  means that  $(\Lambda \otimes A)^{G_i} = (\Lambda)^{G_i} \otimes A$ .

(2) Write  $U = \prod_{j \in J} u_j V(U \cap (\mathbb{A}_F^{\infty})^{\times})$ . We claim that we have  $\text{GL}_2(\mathbb{A}_F^{\infty}) = \prod_{i \in I, j \in J} D^{\times} g_i u_j V(\mathbb{A}_F^{\infty})^{\times}$ , from which the result is immediate. To see this, we need to show that if  $g_i u_j = \delta g_{i'} u_{j'} v z$  then  $i = i'$  and  $j = j'$ .

That  $i = i'$  is immediate from the definition of  $I$ , so we have  $u_{j'} v u_j^{-1} z = g_i^{-1} \delta^{-1} g_i$ . As above, there is some positive integer  $N$  coprime to  $p$  such that  $\delta^N \in F^\times$ , so  $(u_{j'} v u_j^{-1})^N \in (\mathbb{A}_F^\infty)^\times$ . Since  $V$  is normal in  $U$ , we can write  $(u_{j'} v u_j^{-1})^N = (u_{j'} u_j^{-1})^N v'$  for some  $v' \in V$ , so that  $(u_{j'} u_j^{-1})^N \in V(U \cap (\mathbb{A}_F^\infty)^\times)$ . Since  $\#(U/V)$  is a power of  $p$ , we see that in fact  $u_{j'} u_j^{-1} \in V(U \cap (\mathbb{A}_F^\infty)^\times)$ , so that  $j = j'$  by the definition of  $J$ .  $\square$

**5.4. Base change.** We begin by using base change to reduce to a special case. By Facts 4.22 and 4.26, we can replace  $F$  by a solvable totally real extension which is unramified at all primes above  $p$ , and assume that

- $[F : \mathbb{Q}]$  is even.
- $\bar{\rho}$  is unramified outside  $p$ .
- For all primes  $v \nmid p$ , both  $\rho(I_{F_v})$  and  $\rho_0(I_{F_v})$  are unipotent (possibly trivial).
- If  $\rho$  or  $\rho_0$  are ramified at some place  $v \nmid p$ , then  $\bar{\rho}|_{G_{F_v}}$  is trivial, and  $\#k(v) \equiv 1 \pmod{p}$ .
- $\det \rho = \det \rho_0$ . [To see this, note that the assumption that  $\rho, \rho_0$  are crystalline with the same Hodge–Tate weights for all places dividing  $p$  implies that  $\det \rho / \det \rho_0$  is unramified at all places dividing  $p$ . Since we have already assumed that  $\rho(I_{F_v})$  and  $\rho_0(I_{F_v})$  are unipotent for all primes  $v \nmid p$ , we see that the character  $\det \rho / \det \rho_0$  is unramified at all primes, and thus has finite order. Since it is residually trivial, it has  $p$ -power order, and is thus trivial on all complex conjugations; so the extension cut out by its kernel is a finite, abelian, totally real extension unramified at all places dividing  $p$ , as required.]

We will assume from now on that all of these conditions hold. Write  $\chi$  for  $\det \rho = \det \rho_0$ ; then we have  $\chi \varepsilon_p = \chi_{0,l}$  for some algebraic grossencharacter  $\chi_0$ .

From now on, we will assume without further comment that the coefficient field  $L$  is sufficiently large, in the sense that  $L$  contains a primitive  $p$ -th root of unity, and for all  $g \in G_F$ ,  $\mathbb{F}$  contains the eigenvalues of  $\bar{\rho}(g)$ .

**5.5. Patching.** Having used base change to impose the additional conditions of the previous section, we are now in a position to begin the main patching argument.

We let  $D/F$  be a quaternion algebra ramified at exactly the infinite places (which exists by our assumption that  $[F : \mathbb{Q}]$  is even). By the Jacquet–Langlands correspondence, we can and will work with automorphic representations of  $G_D(\mathbb{A}^\infty)$  from now on.

Let  $T_p$  be the set of places of  $F$  lying over  $p$ , let  $T_r$  be the set of primes not lying over  $p$  at which  $\rho$  or  $\rho_0$  is ramified, and let  $T = T_p \amalg T_r$ . If  $v \in T_r$ , write  $\sigma_v$  for a choice of topological generator of  $I_{F_v}/P_{F_v}$ . By our assumptions above, if  $v \in T_r$  then  $\bar{\rho}|_{G_{F_v}}$  is trivial,  $\rho|_{I_{F_v}}, \rho_0|_{I_{F_v}}$  are unipotent, and  $\#k(v) \equiv 1 \pmod{p}$ .

The patching argument will involve the consideration of various finite sets  $Q$  of auxiliary finite places. We will always assume that if  $v \in Q$ , then

- $v \notin T$ ,
- $\#k(v) \equiv 1 \pmod{p}$ , and
- $\bar{\rho}(\text{Frob}_v)$  has distinct eigenvalues, which we denote  $\bar{\alpha}_v$  and  $\bar{\beta}_v$ .

For each set  $Q$  of primes satisfying these conditions, we define deformation problems  $\mathcal{S}_Q = (T \cup Q, \{\mathcal{D}_v\}, \chi)$  and  $\mathcal{S}'_Q = (T \cup Q, \{\mathcal{D}'_v\}, \chi)$  as follows. Let  $\zeta$  be a fixed primitive  $p$ -th root of unity in  $L$ .

- If  $v \in T_p$ , then  $\mathcal{D}_v = \mathcal{D}'_v$  is chosen so that  $R_{\bar{\rho}|_{G_{F_v}}, \chi}^{\square} / I(\mathcal{D}_v) = R_{\bar{\rho}|_{G_{F_v}}, \chi, \text{cr}, \{\text{HT}_{\sigma}(\rho)\}}^{\square}$ .
- If  $v \in Q$ , then  $\mathcal{D}_v = \mathcal{D}'_v$  consists of all lifts of  $\bar{\rho}|_{G_{F_v}}$  with determinant  $\chi$ .
- If  $v \in T_r$ , then  $\mathcal{D}_v$  consists of all lifts of  $\bar{\rho}|_{G_{F_v}}$  with  $\text{char}_{\rho(\sigma_v)}(X) = (X-1)^2$ , while  $\mathcal{D}'_v$  consists of all lifts with  $\text{char}_{\rho(\sigma_v)}(X) = (X-\zeta)(X-\zeta^{-1})$ .

(So, the difference between  $\mathcal{S}_Q$  and  $\mathcal{S}_{\emptyset}$  is that we have allowed our deformations to ramify at places in  $Q$ .) We write

$$R^{\text{loc}} = \widehat{\otimes}_{v \in T, \mathcal{O}} R_{\bar{\rho}|_{G_{F_v}}, \chi}^{\square} / I(\mathcal{D}_v), \quad R^{\text{loc},'} = \widehat{\otimes}_{v \in T, \mathcal{O}} R_{\bar{\rho}|_{G_{F_v}}, \chi}^{\square} / I(\mathcal{D}'_v).$$

Then  $R^{\text{loc}}/\lambda = R^{\text{loc},'}/\lambda$ , because  $\zeta \equiv 1 \pmod{\lambda}$ . In addition, we see from Theorems 3.28 and 3.38 that

- $(R^{\text{loc},'})^{\text{red}}$  is irreducible,  $\mathcal{O}$ -flat, and has Krull dimension  $1 + 3\#T + [F : \mathbb{Q}]$ ,
- $(R^{\text{loc}})^{\text{red}}$  is  $\mathcal{O}$ -flat, equidimensional of Krull dimension  $1 + 3\#T + [F : \mathbb{Q}]$ , and reduction modulo  $\lambda$  gives a bijection between the irreducible components of  $\text{Spec } R^{\text{loc}}$  and those of  $\text{Spec } R^{\text{loc}}/\lambda$ .

We have the global analogues  $R_Q^{\text{univ}} := R_{\bar{\rho}, \mathcal{S}_Q}^{\text{univ}}$ ,  $R_Q^{\text{univ},'} := R_{\bar{\rho}, \mathcal{S}'_Q}^{\text{univ}}$ ,  $R_Q^{\square} := R_{\bar{\rho}, \mathcal{S}_Q}^{\square}$ ,  $R_Q^{\square,'} := R_{\bar{\rho}, \mathcal{S}'_Q}^{\square}$ , and we have  $R_Q^{\text{univ}}/\lambda = R_Q^{\text{univ},'}/\lambda$ ,  $R_Q^{\square}/\lambda = R_Q^{\square,'}/\lambda$ . There are obvious natural maps  $R^{\text{loc}} \rightarrow R_Q^{\square}$ ,  $R^{\text{loc},'} \rightarrow R_Q^{\square,'}$ , and these maps agree after reduction mod  $\lambda$ .

We can and do fix representatives  $\rho_Q^{\text{univ}}, \rho_Q^{\text{univ},'}$  for the universal deformations of  $\bar{\rho}$  over  $R_Q^{\text{univ}}, R_Q^{\text{univ},'}$  respectively, which are compatible with the choices of  $\rho_{\emptyset}^{\text{univ}}, \rho_{\emptyset}^{\text{univ},'}$ , and so that the induced surjections

$$R_Q^{\text{univ}} \twoheadrightarrow R_{\emptyset}^{\text{univ}}, \quad R_Q^{\text{univ},'} \twoheadrightarrow R_{\emptyset}^{\text{univ},'}$$

are identified modulo  $\lambda$ .

Fix a place  $v_0 \in T$ , and set  $\mathcal{J} := \mathcal{O}[[X_{v,i,j}]]_{v \in T, i,j=1,2} / (X_{v_0,1,1})$ . Let  $\mathfrak{a}$  be the ideal of  $\mathcal{J}$  generated by the  $X_{v,i,j}$ . Then our choice of  $\rho_Q^{\text{univ}}$  gives an identification  $R_Q^{\square} \xrightarrow{\sim} R_Q^{\text{univ}} \widehat{\otimes}_{\mathcal{O}} \mathcal{J}$ , corresponding to the universal  $T$ -framed deformation  $(\rho_Q^{\text{univ}}, \{1 + (X_{v,i,j})\}_{v \in T})$ .

Now, by Exercise 3.34, for each place  $v \in Q$  we have an isomorphism  $\rho_Q^{\text{univ}}|_{G_{F_v}} \cong \chi_{\alpha} \oplus \chi_{\beta}$ , where  $\chi_{\alpha}, \chi_{\beta} : G_{F_v} \rightarrow (R_Q^{\text{univ}})^{\times}$ , where  $(\chi_{\alpha} \bmod \mathfrak{m}_{R_Q^{\text{univ}}})(\text{Frob}_v) = \bar{\alpha}_v$ ,  $(\chi_{\beta} \bmod \mathfrak{m}_{R_Q^{\text{univ}}})(\text{Frob}_v) = \bar{\beta}_v$ .

Let  $\Delta_v$  be the maximal  $p$ -power quotient of  $k(v)^\times$ . Then  $\chi_\alpha|_{I_{F_v}}$  factors through the composite

$$I_{F_v} \twoheadrightarrow I_{F_v}/P_{F_v} \twoheadrightarrow k(v)^\times \twoheadrightarrow \Delta_v,$$

and if we write  $\Delta_Q = \prod_{v \in Q} \Delta_v$ ,  $(\prod \chi_\alpha) : \Delta_Q \rightarrow (R_Q^{\text{univ}})^\times$ , then we see that  $(R_Q^{\text{univ}})_{\Delta_Q} = R_\emptyset^{\text{univ}}$ .

The isomorphism  $R_Q^\square \xrightarrow{\sim} R_Q^{\text{univ}} \widehat{\otimes}_{\mathcal{O}} \mathcal{J}$  and the homomorphism  $\Delta_Q \rightarrow (R_Q^{\text{univ}})^\times$  together give a homomorphism  $\mathcal{J}[\Delta_Q] \rightarrow R_Q^\square$ . In the same way, we have a homomorphism  $\mathcal{J}[\Delta_Q] \rightarrow R_Q^{\square'}$ , and again these agree modulo  $\lambda$ . If we write  $\mathfrak{a}_Q := \langle \mathfrak{a}, \delta - 1 \rangle_{\delta \in \Delta_Q} \triangleleft \mathcal{J}[\Delta_Q]$ , then we see that  $R_Q^\square / \mathfrak{a}_Q = R_\emptyset^{\text{univ}}$ , and that  $R_Q^{\square'} / \mathfrak{a}_Q = R_\emptyset^{\text{univ}'}$ , and again these agree modulo  $\lambda$ .

We now examine the spaces of modular forms that we will patch. We have our fixed isomorphism  $\iota : \bar{L} \xrightarrow{\sim} \mathbb{C}$ , and an algebraic grossencharacter  $\chi_0$  such that  $\chi^{\varepsilon_p} = \chi_{0,\iota}$ . Define  $k, \eta$  by  $\text{HT}_\tau(\rho_0) = \{\eta_{\iota\tau}, \eta_{\iota\tau} + k_{\iota\tau} - 1\}$ . We define compact open subgroups  $U_Q = \prod U_{Q,v}$ , where:

- $U_{Q,v} = \text{GL}_2(\mathcal{O}_{F_v})$  if  $v \notin Q \cup T_r$ ,
- $U_{Q,v} = U_0(v) = \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{v} \right\}$  if  $v \in T_r$ , and
- $U_{Q,v} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in U_0(v) \mid a/d \pmod{v} \in k(v)^\times \mapsto 1 \in \Delta_v \right\}$  if  $v \in Q$ .

We let  $\psi : \prod_{v \in T_r} U_{Q,v} \rightarrow \mathcal{O}^\times$  be the trivial character. Similarly, we set  $U'_Q = U_Q$ , and we define  $\psi' : \prod_{v \in T_r} U_{Q,v} \rightarrow \mathcal{O}^\times$  in the following way. For each  $v \in T_r$ , we have a homomorphism  $U_{Q,v} \rightarrow k(v)^\times$  given by sending  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  to  $a/d \pmod{v}$ , and we compose these characters with the characters  $k(v)^\times \rightarrow \mathcal{O}^\times$  sending the image of  $\sigma_v$  to  $\zeta$ , where  $\sigma_v$  is a generator of  $I_{F_v}/P_{F_v}$ .

We obtain spaces of modular forms  $S(U_Q, \mathcal{O})$ ,  $S(U'_Q, \mathcal{O})$  and corresponding Hecke algebras  $\mathbb{T}_{U_Q}$ ,  $\mathbb{T}_{U'_Q}$  (generated by the Hecke operators  $T_v, S_v$  with  $v \notin T \cup Q$ ). Note that  $\psi = \psi' \pmod{\lambda}$ , so we have  $S(U_Q, \mathcal{O})/\lambda = S(U'_Q, \mathcal{O})/\lambda$ . We let  $\mathfrak{m}_\emptyset \triangleleft \mathbb{T}_{U_\emptyset}$  be the ideal generated by  $\lambda$  and the  $\text{tr } \bar{\rho}(\text{Frob}_v) - T_v$ ,  $\det \bar{\rho}(\text{Frob}_v) - \#k(v)S_v$ ,  $v \notin T$ . This is a proper maximal ideal of  $\mathbb{T}_{U_\emptyset}$ , because it is the kernel of the homomorphism  $\mathbb{T}_{U_\emptyset} \rightarrow \mathcal{O} \twoheadrightarrow \mathbb{F}$ , where the map  $\mathbb{T}_{U_\emptyset} \rightarrow \mathcal{O}$  is the one coming from the automorphicity of  $\rho_0$ , sending  $T_v \mapsto \text{tr } \rho_0(\text{Frob}_v)$ ,  $S_v \mapsto \#k(v)^{-1} \det \rho_0(\text{Frob}_v)$ .

By the universal property of  $R_\emptyset^{\text{univ}}$ , we have a surjection  $R_\emptyset^{\text{univ}} \twoheadrightarrow \mathbb{T}_\emptyset := \mathbb{T}_{U_\emptyset, \mathfrak{m}_\emptyset}$ , and a corresponding lifting  $\rho^{\text{mod}} : G_F \rightarrow \text{GL}_2(\mathbb{T}_\emptyset)$  of type  $\mathcal{S}_\emptyset$ . Similarly, we have a surjection  $R_\emptyset^{\text{univ}'}/\lambda \twoheadrightarrow \mathbb{T}'_\emptyset := \mathbb{T}_{U'_\emptyset, \mathfrak{m}_\emptyset}$ . Set  $\mathcal{S}_\emptyset := S(U_\emptyset, \mathcal{O})_{\mathfrak{m}_\emptyset}$ ,  $\mathcal{S}'_\emptyset := S(U'_\emptyset, \mathcal{O})_{\mathfrak{m}_\emptyset}$ . Then the identification  $R_\emptyset^{\text{univ}}/\lambda \cong R_\emptyset^{\text{univ}'}/\lambda$  is compatible with  $\mathcal{S}_\emptyset/\lambda = \mathcal{S}'_\emptyset/\lambda$ .

We claim that in order to show that  $\rho$  is modular, it suffices to show that  $\text{Supp}_{R_\emptyset^{\text{univ}}}(\mathcal{S}_\emptyset) = \text{Spec } R_\emptyset^{\text{univ}}$ . Suppose that this is true; then since  $\mathcal{S}_\emptyset$  is a faithful  $\mathbb{T}_\emptyset$ -module by definition, we see that  $\ker(R_\emptyset^{\text{univ}} \rightarrow \mathbb{T}_\emptyset)$  is nilpotent, so that  $(R_\emptyset^{\text{univ}})^{\text{red}} \xrightarrow{\sim} \mathbb{T}_\emptyset$ . Then  $\rho$  corresponds to some homomorphism  $R_\emptyset^{\text{univ}} \rightarrow \mathcal{O}$ , and thus to a homomorphism  $\mathbb{T}_\emptyset \rightarrow \mathcal{O}$ , and the composite of

this homomorphism with  $\iota : \mathcal{O} \hookrightarrow \mathbb{C}$  corresponds to a cuspidal automorphic representation  $\pi$  of  $G_D(\mathbb{A}^\infty)$  of weight  $(k, \eta)$ , which by construction has the property that  $\rho \cong \rho_{\pi, \iota}$ , as required.

To show that  $\text{Supp}_{R_{\mathcal{O}}^{\text{univ}}}(S_{\mathcal{O}}) = \text{Spec } R_{\mathcal{O}}^{\text{univ}}$ , we will study the above constructions as  $Q$  varies. Let  $\mathfrak{m}_Q \triangleleft \mathbb{T}_{U_Q}$  be the maximal ideal generated by  $\lambda$ , the  $\text{tr } \bar{\rho}(\text{Frob}_v) - T_v$  and  $\det \bar{\rho}(\text{Frob}_v) - \#k(v)S_v$  for  $v \notin T \cup Q$ , and the  $U_{\omega_v} - \bar{\alpha}_v$  for  $v \in Q$ , where

$$U_{\omega_v} = \left[ U_{Q,v} \begin{pmatrix} \omega_v & 0 \\ 0 & 1 \end{pmatrix} U_{Q,v} \right].$$

We write  $S_Q = S_{U_Q} := S(U_Q, \mathcal{O})_{\mathfrak{m}_Q}$ , and  $\mathbb{T}_Q := (\mathbb{T}_{U_Q})_{\mathfrak{m}_Q}$ . We have a homomorphism  $\Delta_Q \rightarrow \text{End}(S_Q)$ , given by sending  $\delta \in \Delta_v$  to  $\begin{pmatrix} \delta & 0 \\ 0 & 1 \end{pmatrix} \in U_0(v)$ . We also have another homomorphism  $\Delta_Q \rightarrow \text{End}(S_Q)$ , given by the composite

$$\Delta_Q \rightarrow R_Q^{\text{univ}} \rightarrow \mathbb{T}_Q \rightarrow \text{End}(S_Q).$$

We now examine the consequences of local-global compatibility at the places in  $Q$ . A homomorphism  $\theta : \mathbb{T}_Q \rightarrow \mathbb{C}$  corresponds to a cuspidal automorphic representation  $\pi$ , and for each  $v \in Q$  the image  $\alpha_v$  of  $U_{\omega_v}$  is such that  $\alpha_v$  is an eigenvalue of  $U_{\omega_v}$  on  $\pi_v^{U_{Q,v}}$ .

It can be checked that since  $\pi_v^{U_{Q,v}} \neq 0$ ,  $\pi_v$  is necessarily a subquotient of  $\chi_1 \times \chi_2$  for some tamely ramified characters  $\chi_1, \chi_2 : F_v^\times \rightarrow \mathbb{C}^\times$ . Then one checks explicitly that

$$(\chi_1 \times \chi_2)^{U_{Q,v}} \cong \mathbb{C}\phi_1 \oplus \mathbb{C}\phi_w,$$

where  $w = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ ,  $\phi_1(1) = \phi_w(w) = 1$ , and  $\text{Supp } \phi_1 = B(F_v)U_{Q,v}$ ,  $\text{Supp } \phi_w = B(F_v)wU_{Q,v}$ .

Further explicit calculation shows that

$$U_{\omega_v}\phi_1 = \#k(v)^{1/2}\chi_1(\pi_v)\phi_1 + X\phi_w$$

for some  $X$ , which is 0 if  $\chi_1/\chi_2$  is ramified, and

$$U_{\omega_v}\phi_w = \#k(v)^{1/2}\chi_2(\pi_v)\phi_w.$$

Note that by local-global compatibility,  $\iota^{-1}(\#k(v)^{1/2}\chi_1(\pi_v))$  and  $\iota^{-1}(\#k(v)^{1/2}\chi_2(\pi_v))$  are the eigenvalues of  $\rho_{\pi, \iota}(\phi_v)$ , so one of them is a lift of  $\bar{\alpha}_v$ , and one is a lift of  $\bar{\beta}_v$ .

It is also easily checked that

$$\begin{pmatrix} \delta & 0 \\ 0 & 1 \end{pmatrix} \phi_1 = \chi_1(\delta)\phi_1, \quad \begin{pmatrix} \delta & 0 \\ 0 & 1 \end{pmatrix} \phi_w = \chi_2(\delta)\phi_w.$$

By local-global compatibility,

$$\rho_{\pi, \iota}|_{W_{F_v}^{\text{ss}}} \cong (\chi_1|\cdot|^{-1/2} \oplus \chi_2|\cdot|^{-1/2}) \circ \text{Art}_{F_v}^{-1} = \chi_\beta \oplus \chi_\alpha,$$

say. Reducing modulo  $\lambda$ , we see that  $\{\bar{\alpha}_v, \bar{\beta}_v\} = \{\#k(v)^{1/2}\iota^{-1}(\bar{\chi}_1(\pi_v)), \#k(v)^{1/2}\iota^{-1}(\bar{\chi}_2(\pi_v))\}$ . As a consequence, we see that  $\chi_1/\chi_2 \neq |\cdot|^{\pm 1}$  (as if this equality held, we would have  $\bar{\alpha}_v/\bar{\beta}_v \equiv$

$\#k(v)^{\pm 1} \equiv 1 \pmod{\lambda}$ , contradicting our assumption that  $\bar{\alpha}_v \neq \bar{\beta}_v$ ). Consequently we have  $\pi_v = \chi_1 \times \chi_2 \cong \chi_2 \times \chi_1$ , so that without loss of generality we have  $\bar{\chi}_1(\pi_v) = \bar{\beta}_v, \bar{\chi}_2(\pi_v) = \bar{\alpha}_v$ .

We see that  $S_Q \otimes_{\mathcal{O},i} \mathbf{C} = \bigoplus_{\pi \otimes_{v \in Q} X_v}$ , where  $X_v$  is the 1-dimensional space where  $U_{\omega_v}$  acts via a lift of  $\bar{\alpha}_v$ . Since this space is spanned by  $\phi_w$ , we see that  $\Delta_v$  acts on  $S_Q$  via  $\chi_2 = \chi_\alpha \circ \text{Art}$ , so that we have proved the following important fact.

5.6. *Fact.* The two homomorphisms  $\Delta_Q \rightarrow \text{End}(S_Q)$  (the other one coming via  $R_Q^{\text{univ}}$ ) are equal.

Let  $U_{Q,0} := \prod_{v \notin Q} U_{Q,v} \prod_{v \in Q} U_0(v)$ . Then  $U_Q$  is a normal subgroup of  $U_{Q,0}$ , and  $U_{Q,0}/U_Q = \Delta_Q$ .

5.7. *Fact.*  $S_Q$  is finite free over  $\mathcal{O}[\Delta_Q]$ .

*Proof.* This is immediate from Proposition 5.3(2).  $\square$

Fix a place  $v \in Q$ . Since  $\bar{\alpha}_v \neq \bar{\beta}_v$ , by Hensel's lemma we may write  $\text{char } \rho_{\mathcal{O}}^{\text{mod}}(\text{Frob}_v) = (X - A_v)(X - B_v)$  for some  $A_v, B_v \in \mathbb{T}_{\mathcal{O}}$  with  $A_v \equiv \bar{\alpha}_v, B_v \equiv \bar{\beta}_v \pmod{\mathfrak{m}_{\mathcal{O}}}$ .

5.8. **Proposition.** *We have an isomorphism  $\prod_{v \in Q} (U_{\omega_v} - B_v) : S_{\mathcal{O}} \xrightarrow{\sim} S(U_{Q,0}, \mathcal{O})_{\mathfrak{m}_Q}$ .*

*Proof.* We claim that it is enough to prove that the map is an isomorphism after tensoring with  $L$ , and an injection after tensoring with  $\mathbb{F}$ . To see this, write  $X := S_{\mathcal{O}}, Y := S(U_{Q,0}, \mathcal{O})_{\mathfrak{m}_Q}$ , and write  $Q$  for the cokernel of the map  $X \rightarrow Y$ . Then  $X, Y$  are finite free  $\mathcal{O}$ -modules, and if the map  $X \otimes L \rightarrow Y \otimes L$  is injective, then so is the map  $X \rightarrow Y$ , so that we have a short exact sequence  $0 \rightarrow X \rightarrow Y \rightarrow Q \rightarrow 0$ . Tensoring with  $L$ , we have  $Q \otimes L = 0$ . Tensoring with  $\mathbb{F}$ , we obtain an exact sequence  $0 \rightarrow Q[\lambda] \rightarrow X \otimes \mathbb{F} \rightarrow Y \otimes \mathbb{F} \rightarrow Q \otimes \mathbb{F} \rightarrow 0$ , so we have  $Q[\lambda] = 0$ . Thus  $Q = 0$ , as required.

In order to check that we have an isomorphism after tensoring with  $L$ , it is enough to check that the induced map  $\prod_{v \in Q} (U_{\omega_v} - B_v) : S_{\mathcal{O}} \otimes_{\mathcal{O},i} \mathbf{C} \rightarrow S(U_{Q,0}, \mathcal{O})_{\mathfrak{m}_Q} \otimes_{\mathcal{O},i} \mathbf{C}$  is an isomorphism. This is easily checked:  $S_{\mathcal{O}} \otimes \mathbf{C} \cong \bigoplus_{\pi \otimes_{v \in Q} (\chi_{1,v} \times \chi_{2,v})^{\text{GL}_2(\mathcal{O}_{F_v})}}$ , and  $(\chi_{1,v} \times \chi_{2,v})^{\text{GL}_2(\mathcal{O}_{F_v})} = \mathbf{C}\phi_0$ , where  $\phi_0$  is as in Exercise 4.7(3). Similarly,  $S(U_{Q,0}, \mathcal{O})_{\mathfrak{m}_Q} \otimes_{\mathcal{O},i} \mathbf{C} = \bigoplus_{\pi \otimes_{v \in Q} M_v$ , where  $M_v$  is the subspace of  $(\chi_{1,v} \times \chi_{2,v})^{U_0(v)}$  on which  $U_{\omega_v}$  acts via a lift of  $\bar{\alpha}_v$ , which is spanned by  $\phi_w$ . Since the natural map  $(\chi_{1,v} \times \chi_{2,v})^{\text{GL}_2(\mathcal{O}_{F_v})} \rightarrow (\chi_{1,v} \times \chi_{2,v})^{U_0(v)}$  sends  $\phi_0 \mapsto \phi_1 + \phi_w$  (as  $\phi_0(1) = \phi_0(w) = 1$ ), the result follows.

It remains to check injectivity after tensoring with  $\mathbb{F}$ . The kernel of the map would be a nonzero finite module for the Artinian local ring  $\mathbb{T}_{\mathcal{O}}/\lambda$ , and would thus have nonzero  $\mathfrak{m}_{\mathcal{O}}$ -torsion, so it suffices to prove that the induced map

$$\prod_{v \in Q} (U_{\omega_v} - B_v) : (S_{\mathcal{O}} \otimes \mathbb{F})[\mathfrak{m}_{\mathcal{O}}] \rightarrow S(U_{Q,0}, \mathcal{O})_{\mathfrak{m}_Q} \otimes \mathbb{F}$$

is an injection. By induction on  $\#Q$ , it suffices to prove this in the case that  $Q = \{v\}$ . Suppose for the sake of contradiction that there is a nonzero  $x \in (S_{\mathcal{O}} \otimes \mathbb{F})[\mathfrak{m}_{\mathcal{O}}]$  with  $(U_{\omega_v} - \bar{\beta}_v)x = 0$ .



Since  $x \in S_\emptyset \otimes \mathbb{F}$ , we also have  $T_v x = (\bar{\alpha}_v + \bar{\beta}_v)x$ , and we will show that these two equations together lead to a contradiction.

Now,  $x$  is just a function  $D^\times \backslash \mathrm{GL}_2(\mathbb{A}_F^\infty) \rightarrow \Lambda \otimes \mathbb{F}$ , on which  $\mathrm{GL}_2(\mathbb{A}_F^\infty)$  acts by right translation. If we make the action of the Hecke operators explicit, we find that there are  $g_i$  such that  $U_v = \coprod_i g_i U_{Q,v}$ , and  $T_v = (\coprod_i g_i \mathrm{GL}_2(\mathcal{O}_{F_v})) \coprod \begin{pmatrix} 1 & 0 \\ 0 & \pi_v \end{pmatrix} \mathrm{GL}_2(\mathcal{O}_{F_v})$ , so that we have  $\begin{pmatrix} 1 & 0 \\ 0 & \pi_v \end{pmatrix} x = T_v x - U_{\omega_v} x = \bar{\alpha}_v x$ . Then  $\begin{pmatrix} \pi_v & 0 \\ 0 & 1 \end{pmatrix} x = w \begin{pmatrix} 1 & 0 \\ 0 & \pi_v \end{pmatrix} w x = \bar{\alpha}_v x$ , and  $U_{\omega_v} x = \sum_{a \in k(v)} \begin{pmatrix} \pi_v & a \\ 0 & 1 \end{pmatrix} x = \sum_{a \in k(v)} \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \pi_v & 0 \\ 0 & 1 \end{pmatrix} x = \#k(v) \bar{\alpha}_v x = \bar{\alpha}_v x$ . But  $U_{\omega_v} x = \bar{\beta}_v x$ , so  $\bar{\alpha}_v = \bar{\beta}_v$ , a contradiction.  $\square$

Set  $S_Q^\square := S_Q \otimes_{R_Q^{\mathrm{univ}}} R_Q^\square$ . Then we have  $S_Q^\square / \mathfrak{a}_Q = S(U_{Q,0}, \mathcal{O})_{\mathfrak{m}_Q} \xrightarrow{\sim} S_\emptyset$ , compatibly with the isomorphism  $R_Q^\square / \mathfrak{a}_Q \xrightarrow{\sim} R_\emptyset^{\mathrm{univ}}$ . Also,  $S_Q^\square$  is finite free over  $\mathcal{J}[\Delta_Q]$ .

We now return to the Galois side. By Proposition 3.24, we can and do choose a presentation

$$R^{\mathrm{loc}}[[x_1, \dots, x_{h_Q}]] \twoheadrightarrow R_Q^{\square T},$$

where  $h_Q = \#T + \#Q - 1 - [F : \mathbb{Q}] + \dim_{\mathbb{F}} H_Q^1(G_{F,T}, (\mathrm{ad}^0 \bar{\rho})(1))$ , and  $H_Q^1(G_{F,T}, (\mathrm{ad}^0 \bar{\rho})(1)) = \ker(H^1(G_{F,T}, (\mathrm{ad}^0 \bar{\rho})(1)) \rightarrow \bigoplus_{v \in Q} H^1(G_{k(v)}, (\mathrm{ad}^0 \bar{\rho})(1)))$ .

The following result will provide us with the sets  $Q$  that we will use.

**5.9. Proposition.** *Let  $r = \max(\dim H^1(G_{F,T}, (\mathrm{ad}^0 \bar{\rho})(1)), 1 + [F : \mathbb{Q}] - \#T)$ . For each  $N \geq 1$ , there exists a set  $Q_N$  of primes of  $F$  such that*

- $Q_N \cap T = \emptyset$ .
- If  $v \in Q_N$ , then  $\bar{\rho}(\mathrm{Frob}_v)$  has distinct eigenvalues  $\bar{\alpha}_v \neq \bar{\beta}_v$ .
- If  $v \in Q_N$ , then  $\#k(v) \equiv 1 \pmod{p^N}$ .
- $\#Q_N = r$ .
- $R_{Q_N}^\square$  (respectively  $R_{Q_N}^{\square'}$ ) is topologically generated over  $R^{\mathrm{loc}}$  (respectively  $R^{\mathrm{loc}'}$ ) by  $\#T - 1 - [F : \mathbb{Q}] + r$  elements.

*Proof.* The last condition may be replaced by

- $H_{Q_N}^1(G_{F,T}, (\mathrm{ad}^0 \bar{\rho})(1)) = (0)$ .

Therefore, it is enough to show that for each  $0 \neq [\phi] \in H^1(G_{F,T}, (\mathrm{ad}^0 \bar{\rho})(1))$ , there are infinitely many  $v \notin T$  such that

- $\#k(v) \equiv 1 \pmod{p^N}$ .
- $\bar{\rho}(\mathrm{Frob}_v)$  has distinct eigenvalues  $\bar{\alpha}_v, \bar{\beta}_v$ .
- $\mathrm{Res}[\phi] \in H^1(G_{k(v)}, (\mathrm{ad}^0 \bar{\rho})(1))$  is nonzero.

(This then gives us some set of primes  $Q$  with the given properties, except that  $\#Q$  may be too large; but then we can pass to a subset of cardinality  $r$ , while maintaining the injectivity of the map  $H^1(G_{F,T}, (\text{ad}^0 \bar{\rho})(1)) \rightarrow \bigoplus_{v \in Q} H^1(G_{k(v)}, (\text{ad}^0 \bar{\rho})(1))$ .)

We will use the Chebotarev density theorem to do this; note that the condition that  $\#k(v) \equiv 1 \pmod{p^N}$  is equivalent to  $v$  splitting completely in  $F(\zeta_{p^N})$ , and the condition that  $\bar{\rho}(\text{Frob}_v)$  has distinct eigenvalues is equivalent to asking that  $\text{ad}^0 \bar{\rho}(\text{Frob}_v)$  has an eigenvalue not equal to 1.

Set  $E = \bar{F}^{\ker \text{ad}^0 \bar{\rho}}(\zeta_{p^N})$ . We claim that we have  $H^1(\text{Gal}(E/F), (\text{ad}^0 \bar{\rho})(1)) = (0)$ . In order to see this, we claim firstly that  $\zeta_p \notin \bar{F}^{\ker \text{ad}^0 \bar{\rho}}$ . This follows from the classification of finite subgroups of  $\text{PGL}_2(\bar{\mathbb{F}}_p)$ : we have assumed that  $\text{Im} \bar{\rho} \supseteq \text{SL}_2(\mathbb{F}_p)$ , and this implies that  $\text{Im} \text{ad}^0 \bar{\rho} = \text{PGL}_2(\mathbb{F}_{p^s})$  or  $\text{PSL}_2(\mathbb{F}_{p^s})$  for some  $s$ , and in particular  $(\text{Im} \text{ad}^0 \bar{\rho})^{\text{ab}}$  is trivial or cyclic of order 2. Since  $p \geq 5$  and  $p$  is unramified in  $F$ , we have  $[F(\zeta_p) : F] \geq 4$ , so  $\zeta_p \notin \bar{F}^{\ker \text{ad}^0 \bar{\rho}}$ , as claimed.

The extension  $E/\bar{F}^{\ker \text{ad}^0 \bar{\rho}}$  is abelian, and we let  $E_0$  be the intermediate field such that  $\text{Gal}(E/E_0)$  has order prime to  $p$ , while  $\text{Gal}(E_0/\bar{F}^{\ker \text{ad}^0 \bar{\rho}})$  has  $p$ -power order. Write  $\Gamma_1 = \text{Gal}(E_0/F)$ ,  $\Gamma_2 = \text{Gal}(E/E_0)$ . Then the inflation-restriction exact sequence is in part

$$(0) \rightarrow H^1(\Gamma_1, (\text{ad}^0 \bar{\rho})(1)^{\Gamma_2}) \rightarrow H^1(\text{Gal}(E/F), (\text{ad}^0 \bar{\rho})(1)) \rightarrow H^1(\Gamma_2, (\text{ad}^0 \bar{\rho})(1))^{\Gamma_1},$$

so in order to show that  $H^1(\text{Gal}(E/F), (\text{ad}^0 \bar{\rho})(1)) = (0)$ , it suffices to prove that  $H^1(\Gamma_1, (\text{ad}^0 \bar{\rho})(1)^{\Gamma_2}) = H^1(\Gamma_2, (\text{ad}^0 \bar{\rho})(1))^{\Gamma_1} = (0)$ .

In fact, we claim that  $(\text{ad}^0 \bar{\rho})(1)^{\Gamma_2}$  and  $H^1(\Gamma_2, (\text{ad}^0 \bar{\rho})(1))$  both vanish. For the first of these, note that  $\Gamma_2$  acts trivially on  $\text{ad}^0 \bar{\rho}$  (since  $E_0$  contains  $\bar{F}^{\ker \text{ad}^0 \bar{\rho}}$ ), but that  $\zeta_p \notin E_0$  (as  $[E_0 : \bar{F}^{\ker \text{ad}^0 \bar{\rho}}]$  is a power of  $p$ ). For the second term, note that  $\Gamma_2$  has prime-to- $p$  order.

Suppose that  $\#k(v) \equiv 1 \pmod{p}$ , and that  $\bar{\rho}(\text{Frob}_v) = \begin{pmatrix} \bar{\alpha}_v & 0 \\ 0 & \bar{\beta}_v \end{pmatrix}$ . Then  $\text{ad}^0 \bar{\rho}$  has the basis  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$  of eigenvectors for  $\text{Frob}_v$ , with eigenvalues  $1, \bar{\alpha}_v/\bar{\beta}_v, \bar{\beta}_v/\bar{\alpha}_v$  respectively. Consequently, we see that there is an isomorphism  $H^1(G_{k(v)}, (\text{ad}^0 \bar{\rho})(1)) \cong \mathbb{F}$  (since in general for a (pro)cyclic group, the first cohomology is given by passage to coinvariants), which we can write explicitly as  $[\phi] \mapsto \pi_v \circ \phi(\text{Frob}_v) \circ i_v$ , where  $i_v$  is the injection of  $\mathbb{F}$  into the  $\bar{\alpha}_v$ -eigenspace of  $\text{Frob}_v$ , and  $\pi_v$  is the  $\text{Frob}_v$ -equivariant projection onto that subspace.

Let  $\sigma_0$  be an element of  $\text{Gal}(E/F)$  such that

- $\sigma_0(\zeta_{p^N}) = \zeta_{p^N}$ .
- $\bar{\rho}(\sigma_0)$  has distinct eigenvalues  $\bar{\alpha}, \bar{\beta}$ .

(To see that such a  $\sigma_0$  exists, note that  $\text{Gal}(\bar{F}^{\ker \bar{\rho}}/F(\zeta_{p^N}) \cap \bar{F}^{\ker \bar{\rho}})$  contains  $\text{PSL}_2(\mathbb{F}_p)$ , and so we can choose  $\sigma_0$  so that its image in this group is an element whose adjoint has an eigenvalue other than 1.) Let  $\tilde{E}/E$  be the extension cut out by all the  $[\phi] \in H^1(G_{F,T}, (\text{ad}^0 \bar{\rho})(1))$ . In order to

complete the proof, it suffices to show that we can choose some  $\sigma \in \text{Gal}(\tilde{E}/F)$  with  $\sigma|_E = \sigma_0$ , and such that in the notation above, we have  $\pi_{\sigma_0} \circ \phi(\sigma) \circ i_{\sigma_0} \neq 0$ , because we can then choose  $v$  to have  $\text{Frob}_v = \sigma$  by the Chebotarev density theorem.

To this end, choose any  $\tilde{\sigma}_0 \in \text{Gal}(\tilde{E}/F)$  with  $\tilde{\sigma}_0|_E = \sigma_0$ . If  $\tilde{\sigma}_0$  does not work, then we have  $\pi_{\sigma_0} \circ \phi(\tilde{\sigma}_0) \circ i_{\sigma_0} = 0$ . In this case, take  $\sigma = \sigma_1 \tilde{\sigma}_0$  for some  $\sigma_1 \in \text{Gal}(\tilde{E}/E)$ . Then  $\phi(\sigma) = \phi(\sigma_1 \tilde{\sigma}_0) = \phi(\sigma_1) + \sigma_1 \phi(\tilde{\sigma}_0) = \phi(\sigma_1) + \phi(\tilde{\sigma}_0)$ , so  $\pi_{\sigma_0} \circ \phi(\sigma) \circ i_{\sigma_0} = \pi_{\sigma_0} \circ \phi(\sigma_1) \circ i_{\sigma_0}$ .

Note that  $\phi(\text{Gal}(\tilde{E}/E))$  is a  $\text{Gal}(E/F)$ -invariant subset of  $\text{ad}^0 \bar{\rho}$ , which is an irreducible  $\text{Gal}(E/F)$ -module, since the image of  $\bar{\rho}$  contains  $\text{SL}_2(\mathbb{F}_p)$ . Thus the  $\mathbb{F}$ -span of  $\phi(\text{Gal}(\tilde{E}/E))$  is all of  $\text{ad}^0 \bar{\rho}$ , from which it is immediate that we can choose  $\sigma_1$  so that  $\pi_{\sigma_0} \circ \phi(\sigma_1) \circ i_{\sigma_0} \neq 0$ .  $\square$

We are now surprisingly close to proving the main theorem! Write  $h := \#T - 1 - [F : \mathbb{Q}] + r$ , and  $R_\infty := R^{\text{loc}}[[x_1, \dots, x_h]]$ . For each set  $Q_N$  as above, choose a surjection  $R_\infty \twoheadrightarrow R_{Q_N}^\square$ . Let  $\mathcal{J}_\infty := \mathcal{J}[[y_1, \dots, y_r]]$ . Choose a surjection  $\mathcal{J}_\infty \twoheadrightarrow \mathcal{J}[\Delta_{Q_N}]$ , given by writing  $Q_N = \{v_1, \dots, v_r\}$  and mapping  $y_i$  to  $(\gamma_i - 1)$ , where  $\gamma_i$  is a generator of  $\Delta_{v_i}$ . Choose a homomorphism  $\mathcal{J}_\infty \rightarrow R_\infty$  so that the composites  $\mathcal{J}_\infty \rightarrow R_\infty \twoheadrightarrow R_{Q_N}^\square$  and  $\mathcal{J}_\infty \rightarrow \mathcal{J}[\Delta_{Q_N}] \rightarrow R_{Q_N}^\square$  agree, and write  $\mathfrak{a}_\infty := (\mathfrak{a}, y_1, \dots, y_r)$ . Then  $S_{Q_N}^\square / \mathfrak{a}_\infty = S_\emptyset$ ,  $R_{Q_N}^\square / \mathfrak{a}_\infty = R_\emptyset^{\text{univ}}$ .

Write  $\mathfrak{b}_N := \ker(\mathcal{J}_\infty \rightarrow \mathcal{J}[\Delta_{Q_N}])$ , so that  $S_{Q_N}^\square$  is finite free over  $\mathcal{J}_\infty / \mathfrak{b}_N$ . Since all the elements of  $Q_N$  are congruent to 1 modulo  $p^N$ , we see that  $\mathfrak{b}_N \subseteq ((1 + y_1)^{p^N} - 1, \dots, (1 + y_r)^{p^N} - 1)$ .

We can and do choose the same data for  $R^{\text{loc}'}$ , in such a way that the two sets of data are compatible modulo  $\lambda$ .

Now choose open ideals  $\mathfrak{c}_N \triangleleft \mathcal{J}_\infty$  such that

- $\mathfrak{c}_N \cap \mathcal{O} = (\lambda^N)$ .
- $\mathfrak{c}_N \supseteq \mathfrak{b}_N$ .
- $\mathfrak{c}_N \supseteq \mathfrak{c}_{N+1}$ .
- $\bigcap_N \mathfrak{c}_N = (0)$ .

(For example, we could take  $\mathfrak{c}_N = ((1 + X_{v_i,j})^{p^N} - 1, (1 + y_i)^{p^N} - 1, \lambda^N)$ .) Note that since  $\mathfrak{c}_N \supseteq \mathfrak{b}_N$ ,  $S_{Q_N}^\square / \mathfrak{c}_N$  is finite free over  $\mathcal{J}_\infty / \mathfrak{c}_N$ . Also choose open ideals  $\mathfrak{d}_N \triangleleft R_\emptyset^{\text{univ}}$  such that

- $\mathfrak{d}_N \subseteq \ker(R_\emptyset^{\text{univ}} \rightarrow \text{End}(S_\emptyset / \lambda^N))$ .
- $\mathfrak{d}_N \supseteq \mathfrak{d}_{N+1}$ .
- $\bigcap_N \mathfrak{d}_N = (0)$ .

If  $M \geq N$ , write  $S_{M,N} = S_{\mathcal{O}_M}^{\square} / \mathfrak{c}_N$ , so that  $S_{M,N}$  is finite free over  $\mathcal{J}_{\infty} / \mathfrak{c}_N$  of rank equal to the  $\mathcal{O}$ -rank of  $S_{\emptyset}$ ; indeed  $S_{M,N} / \mathfrak{a}_{\infty} \xrightarrow{\sim} S_{\emptyset} / \lambda^N$ . Then we have a commutative diagram

$$\begin{array}{ccc} \mathcal{J}_{\infty} & \longrightarrow & R_{\infty} \longrightarrow \twoheadrightarrow R_{\emptyset}^{\text{univ}} / \mathfrak{d}_N \\ & & \downarrow \quad \quad \downarrow \\ & & S_{M,N} \longrightarrow \twoheadrightarrow S_{\emptyset} / \mathfrak{d}_N \end{array}$$

where  $S_{M,N}$ ,  $S_{\emptyset} / \mathfrak{d}_N$  and  $R_{\emptyset}^{\text{univ}} / \mathfrak{d}_N$  all have finite cardinality. Because of this finiteness, we see that there is an infinite subsequence of pairs  $(M_i, N_i)$  such that  $M_{i+1} > M_i$ ,  $N_{i+1} > N_i$ , and the induced diagram

$$\begin{array}{ccc} \mathcal{J}_{\infty} & \longrightarrow & R_{\infty} \longrightarrow \twoheadrightarrow R_{\emptyset}^{\text{univ}} / \mathfrak{d}_{N_i} \\ & & \downarrow \quad \quad \downarrow \\ & & S_{M_{i+1}, N_{i+1}} / \mathfrak{c}_{N_i} \longrightarrow \twoheadrightarrow S_{\emptyset} / \mathfrak{d}_{N_i} \end{array}$$

is isomorphic to the diagram for  $(M_i, N_i)$ .

Then we can take the projective limit over this subsequence, to obtain a commutative diagram

$$\begin{array}{ccc} \mathcal{J}_{\infty} & \longrightarrow & R_{\infty} \longrightarrow \twoheadrightarrow R_{\emptyset}^{\text{univ}} \\ & & \downarrow \quad \quad \downarrow \\ & & S_{\infty} \longrightarrow \twoheadrightarrow S_{\emptyset} \end{array}$$

where  $S_{\infty}$  is finite free over  $\mathcal{J}_{\infty}$ . Furthermore, we can simultaneously carry out the same construction in the  $'$  world, compatibly with this picture modulo  $\lambda$ .

This is the key picture, and the theorem will now follow from it by purely commutative algebra arguments. We have  $\dim R_{\infty} = \dim R'_{\infty} = \dim \mathcal{J}_{\infty} = 4\#T + r$ , and since  $S_{\infty}, S'_{\infty}$  are finite free over the power series ring  $\mathcal{J}_{\infty}$ , we have  $\text{depth}_{\mathcal{J}_{\infty}}(S_{\infty}) = \text{depth}_{\mathcal{J}_{\infty}}(S'_{\infty}) = 4\#T + r$ . Since the action of  $\mathcal{J}_{\infty}$  on  $S_{\infty}$  factors through  $R_{\infty}$ , we see that  $\text{depth}_{R_{\infty}}(S_{\infty}) \geq 4\#T + r$ , and similarly  $\text{depth}_{R'_{\infty}}(S'_{\infty}) \geq 4\#T + r$ . Now, if  $\mathcal{P} \triangleleft R'_{\infty}$  is a minimal prime in the support of  $S'_{\infty}$ , then we see that

$$4\#T + r = \dim R'_{\infty} \geq \dim R'_{\infty} / \mathcal{P} \geq \text{depth}_{R'_{\infty}} S'_{\infty} \geq 4\#T + r,$$

so equality holds throughout, and  $\mathcal{P}$  is a minimal prime of  $R'_{\infty}$ . But  $R'_{\infty}$  has a unique minimal prime, so in fact  $\text{Supp}_{R'_{\infty}}(S'_{\infty}) = \text{Spec } R'_{\infty}$ .

By the same argument, we see that  $\text{Supp}_{R_{\infty}}(S_{\infty})$  is a union of irreducible components of  $\text{Spec } R_{\infty}$ . We will show that it is all of  $\text{Spec } R_{\infty}$  by reducing modulo  $\lambda$  and comparing with the situation for  $S'_{\infty}$ .

To this end, note that since  $\text{Supp}_{R'_\infty}(S'_\infty) = \text{Spec } R'_\infty$ , we certainly have  $\text{Supp}_{R'_\infty/\lambda}(S'_\infty/\lambda) = \text{Spec } R'_\infty/\lambda$ . By the compatibility between the two pictures, this means that  $\text{Supp}_{R_\infty/\lambda}(S_\infty/\lambda) = \text{Spec } R_\infty/\lambda$ . Thus  $\text{Supp}_{R_\infty}(S_\infty)$  is a union of irreducible components of  $\text{Spec } R_\infty$ , which contains the entirety of  $\text{Spec } R_\infty/\lambda$ . Since the irreducible components of  $\text{Spec } R_\infty/\lambda$  are in bijection with the irreducible components of  $\text{Spec } R_\infty$ , this implies that  $\text{Supp}_{R_\infty}(S_\infty) = \text{Spec } R_\infty$ . Then  $\text{Supp}_{R_\infty/\mathfrak{a}_\infty}(S_\infty/\mathfrak{a}_\infty) = R_\infty/\mathfrak{a}_\infty$ , i.e.  $\text{Supp}_{R_\infty^{\text{univ}}} S_\infty = R_\infty^{\text{univ}}$ , which is what we wanted to prove.

## 6. LEVEL RAISING AND LOWERING

**6.1. Level raising and lowering: the Khare–Wintenberger method.** So far, our arguments have been directed towards proving the modularity of Galois representations. It turns out that they can also be applied to the internal study of modular forms, by producing congruences between modular forms. Of course, these congruences can also be stated as congruences between the associated Galois representations, and indeed are often best stated in these terms.

The kind of congruences that we will be particularly concerned with are of the following kind: given an automorphic representation  $\pi$  and the associated  $p$ -adic Galois representation  $\rho_{\pi,\iota}$ , we would like to find another automorphic representation  $\pi'$  such that  $\bar{\rho}_{\pi',\iota} \equiv \bar{\rho}_{\pi,\iota}$  and such that  $\pi'$  has different local properties to  $\pi$ . What do we mean by “different local properties”? There are several things we could mean by this, but for the purposes of the present discussion, we will mean that at some primes  $v \nmid p$ ,  $\pi_v$  and  $\pi'_v$  have different conductors. If the conductor of  $\pi'_v$  is greater than that of  $\pi_v$ , we are considering “level raising”, and if the conductor of  $\pi'_v$  is less than that of  $\pi_v$ , the problem is called “level lowering”.

Both of these problems long predate modularity lifting theorems, and indeed results on both were ingredients in the original Taylor–Wiles arguments. However, the most general results now known about these problems come from modularity lifting theorems, and indeed there is now a fairly complete theory, even for  $n$ -dimensional representations. We will now sketch the argument; one of the first exercises is to fill in the details.

It is possible to rephrase the problem in a fashion that makes the connection to modularity lifting theorems more apparent. Suppose that we have a Galois representation  $\rho$  which satisfies the hypotheses of our main theorem with  $\rho_0 = \rho_{\pi,\iota}$ . Then  $\rho$  is modular, so  $\rho \cong \rho_{\pi',\iota}$  for some  $\pi'$ . Then arranging that  $\pi'_v$  has the required conductor is (by the local Langlands correspondence) simply a question of arranging that  $\rho|_{G_{F_v}}$  corresponds to a point of a deformation ring of a particular type. In fact, we can even ask for refinements, where we specify precisely which component of the local deformation ring we should be on.

So, modularity lifting theorems reduce the problem of level raising and level lowering to that of producing a particular Galois representation. At first sight, this may not be an obvious simplification, but we can now rephrase the problem in terms of universal Galois deformation

rings. Indeed, the representation  $\rho$  corresponds to a  $\overline{\mathbb{Q}}_p$ -point on an appropriate Galois deformation ring  $R_S^{\text{univ}}$  (given by choosing an inertial type at finitely many primes  $v \nmid p$ , and by the Fontaine–Laffaille condition at places  $v|p$ ), so it is enough to check that  $R_S^{\text{univ}}$  has  $\overline{\mathbb{Q}}_p$ -points.

You might hope that this could be accomplished by purely commutative algebra arguments, but this seems to be hard. However, we can make a start using Proposition 3.24; under the Taylor–Wiles assumption that  $\bar{\rho}|_{G_F(\zeta_p)}$  is irreducible, we see that  $\dim R_S^{\text{univ}} \geq 1$ . On its own, this doesn't give us anything, because  $R_S^{\text{univ}}$  could be a power series ring over  $\mathbb{F}$ . However, we claim that it is also the case that  $R_S^{\text{univ}}$  is finite over  $\mathcal{O}$ ; this is enough, by (for example) Proposition 2.2 of [KW09].

In order to show that  $R_S^{\text{univ}}$  is finite over  $\mathcal{O}$ , we return to the world of modularity lifting theorems. Indeed, an output of the modularity lifting theorem we proved is that  $R_{\mathcal{O}}^{\text{univ}}$  is a finite  $\mathcal{O}$ -module. To see this, note that we have identified  $(R_{\mathcal{O}}^{\text{univ}})^{\text{red}}$  with a certain Hecke algebra, which is certainly finite over  $\mathcal{O}$  (as it is a subalgebra of the endomorphism ring of a finite  $\mathcal{O}$ -module). Then  $R_{\mathcal{O}}^{\text{univ}}/\mathfrak{m}_{R_{\mathcal{O}}^{\text{univ}}}$  is 0-dimensional and Noetherian, hence Artinian, and it follows from the topological version of Nakayama's lemma that  $R_{\mathcal{O}}^{\text{univ}}$  is finite over  $\mathcal{O}$ .

So, if we are in a situation where  $R_S^{\text{univ}}$  satisfies the hypotheses that we imposed on  $R_{\mathcal{O}}^{\text{univ}}$ , then we are done. Unfortunately, it is not always the case that those hypotheses are satisfied; indeed, a crucial part of the argument was to use base change to put ourselves in the situation that all of the primes at which  $\rho$  or  $\rho_0$  were ramified satisfied some rather restrictive hypotheses. This might seem to be a limitation, but we are rescued by Proposition 3.26. This shows that in order to show that  $R_S^{\text{univ}}$  is finite over  $\mathcal{O}$ , it is enough to show that  $R_{S'}^{\text{univ}}$  is finite over  $\mathcal{O}$ , where  $S'$  is the base change of  $S$  (which is defined in the obvious way); so if we make the base change that we made at the start of the proof of the main theorem, we can write  $R_{S'}^{\text{univ}} = R_{\mathcal{O}}^{\text{univ}}$ , and we are done.

## 7. THE PROJECT (ARIZONA WINTER SCHOOL MARCH 2013)

The first three items below should all be achievable in the project; the later items are more speculative, and will require some new ideas.

- (1) Generalise the main result of [DT94] to Hilbert modular forms, under the usual Taylor–Wiles condition (i.e. that if  $\bar{\rho} : G_F \rightarrow \text{GL}_2(\overline{\mathbb{F}}_p)$  is the representation under consideration, then  $\bar{\rho}|_{G_F(\zeta_p)}$  is irreducible). (This is mostly a matter of expanding the sketch given in Section 6.1.)
- (2) Further generalise this result to allow Hilbert modular forms of weight other than parallel weight two.
- (3) Prove the corresponding results for automorphic forms on quaternion algebras.
- (4) To what extent can these results be combined with results on the weight part of Serre's conjecture, to allow finer control on the representations at  $p$ ?
- (5) Is it possible to relax the assumption that the Taylor–Wiles condition holds?

- (6) What about the case  $p = 2$ ? (Note that the case  $p = 3$  is discussed in section 8.1 below.)

There are also some interesting questions relating to topics that have not been covered in these notes, but which should be accessible without learning huge amounts of new material.

- (7) What are the most general statements about multiplicity one for mod  $p$  automorphic forms on quaternion algebras that can be proved via modularity lifting theorems, using the method of [Dia97] (cf. the recent preprint [BD12] for some results for totally real fields).

There are analogous automorphy lifting theorems available for rank 2 unitary groups (indeed, for rank  $n$  unitary groups), and some results on level raising and lowering are available (cf. Theorem 3.1.2 of [BLGG13]), but they all involve an assumption that should be unnecessary, namely that the Galois representations are only ramified at primes that split in the splitting field of the unitary group.

- (8) Remove this condition, by developing the deformation theory at nonsplit primes.

## 8. GENERALISATIONS

**8.1. Relaxing the hypotheses.** The hypotheses on our main theorem are not optimal. We will now briefly indicate the “easy” relaxations of the assumptions that could be made, and discuss the generalisations that are possible with (a lot) more work.

Firstly, it is possible to relax the assumption that  $p \geq 5$ , and that  $\text{Im } \bar{\rho} \supseteq \text{SL}_2(\mathbb{F}_p)$ . These assumptions cannot be completely removed, but they can be considerably relaxed. The case  $p = 2$  is harder in several ways (and indeed there are no automorphy lifting theorems available for  $\text{GL}_n$  and  $p = 2$  with  $n > 2$ ), and I will say nothing more about it, except that important theorems have been proved in this case, for example the results of [Kis09b] which completed the proof of Serre’s conjecture.

On the other hand, the case  $p = 3$  presents no real difficulties, and we can also allow the image of  $\bar{\rho}$  to be considerably smaller. Firstly, we consider the case  $p = 3$ . The first place that we assumed that  $p > 3$  was in the proof of Proposition 5.3; this argument could also break down for cases when  $p > 3$  if we allowed  $p$  to ramify in  $F$ , which in general we would like to do. Fortunately, there is a simple solution to this problem, which is to introduce an auxiliary prime  $v$  to the level, to make it sufficiently small that the finite groups  $G_i$  have order prime to  $p$ . This prime is chosen in such a way that all deformations of  $\bar{\rho}|_{G_{\bar{F}_v}}$  are automatically unramified, so none of the global Galois deformation rings that we work with are changed when we relax the conditions at  $v$ . The existence of an appropriate  $v$  follows from the Chebotarev density theorem and some elementary group theory; see Lemma 4.11 of [DDT97] and the discussion immediately preceding it.

We now consider the possibility of relaxing the assumption that  $\text{Im } \bar{\rho} \supseteq \text{SL}_2(\mathbb{F}_p)$ . We should certainly assume that  $\bar{\rho}$  is absolutely irreducible, because otherwise many of our constructions

don't even make sense; we always had to assume this in constructing universal deformation rings, in constructing the universal modular deformation, and so on. (Similar theorems have been proved in the case that  $\bar{\rho}$  is reducible, cf. [SW99], but the arguments are considerably more involved, and at present involve a number of serious additional hypotheses, in particular ordinarity.) Examining the arguments made above, we see that the main uses of the assumption that  $\text{Im } \bar{\rho} \supseteq \text{SL}_2(\mathbb{F}_p)$  is in the proof of Proposition 5.9. Looking more closely at the proof, the key assumption is really that  $\bar{\rho}|_{G_{F(\zeta_p)}}$  is absolutely irreducible; this is known as the “Taylor–Wiles assumption”. (Note that by elementary group theory, this is equivalent to the absolute irreducibility of  $\bar{\rho}|_{G_K}$ , where  $K/F$  is the unique quadratic subextension of  $F(\zeta_p)/F$ ; in particular, over  $\mathbb{Q}$  the condition is equivalent to the absolute irreducibility of  $\bar{\rho}|_{G_{\mathbb{Q}(\sqrt{(-1)^{(p-1)/2}p})}}$ , which is how the condition is stated in the original papers.)

Unfortunately this condition isn't quite enough in complete generality, but it comes very close; the only exception is certain cases when  $p = 5$ ,  $F$  contains  $\mathbb{Q}(\sqrt{5})$ , and the projective image of  $\bar{\rho}$  is  $\text{PGL}_2(\mathbb{F}_5)$ . See [Kis09c, (3.2.3)] for the definitive statement. Without assuming that  $\bar{\rho}|_{G_{F(\zeta_p)}}$  is absolutely irreducible, the argument runs into trouble at this point, and the only results are those of [SW01] in the ordinary case, which use similar arguments to those of [SW99].

The other conditions that we could hope to relax are the assumptions on  $\rho|_{G_{F_v}}, \rho_0|_{G_{F_v}}$  at places  $v|p$ . We've hardly discussed where some of these assumptions come from, as we swept most issues with  $p$ -adic Hodge theory under the carpet. There are essentially two problems here. One is that we have assumed that  $p$  is unramified in  $F$ , that the Galois representations are crystalline, and that the gaps between the Hodge–Tate weights are “small”; this is the Fontaine–Laffaille condition. There is also the assumption that  $\rho, \rho_0$  have the same Hodge–Tate weights.

Both conditions can be considerably (although by no means completely) relaxed. For the first restriction, the fundamental difficulty is in understanding the local deformation rings, which are no longer smooth in general. In the Fontaine–Laffaille case, the integral  $p$ -adic Hodge theory is very simple, and amenable to direct computation. In any greater generality, while we have suitable integral  $p$ -adic Hodge theory, it seems hopeless to make general computations (essentially the only computations that have been made are for certain tamely potentially Barsotti–Tate representations; that is, for potentially crystalline representations, where the gaps between the Hodge–Tate weights are all equal to 1, and the representations become crystalline over a tamely ramified extension). Furthermore, while very little is known about these local deformation rings, it seems very likely that they have many irreducible components in general.

This is a problem, because if we run through the arguments above in a more general context (assuming for the sake of exposition that  $T$  consists of just a single place lying over  $p$ ), we will find that the support of  $R_\emptyset$  will be a union of components of the local deformation ring, and we need to show that it is supported on every component. This is an analogous problem to the



one that we faced at the places in  $T_r$ , but it seems hard to make a similar argument, because in general we have no way of explicitly computing the deformation rings. At present, there are three methods for getting around this problem, none of which is strictly stronger than the other.

The first method is that of [Kis09c], which applies only in the potentially Barsotti–Tate case, but which proves essentially complete results in this case. In this case (which was also the case considered in the original work of Taylor–Wiles on the modularity of elliptic curves, and the subsequent work of Breuil–Conrad–Diamond–Taylor) the local deformation ring can be reinterpreted in terms of  $p$ -divisible groups and finite flat group schemes. After making a base change to the Barsotti–Tate (i.e. crystalline case), the deformations being considered are those which are the generic fibres of finite flat group schemes. In a highly ramified situation, a representation can have many finite flat models, and Kisin constructs a moduli space of these models, which is better behaved than the Galois deformation ring. This space can be studied in a relatively explicit fashion, and (possibly after a further base change) it is possible to check that the Galois deformation ring has at most two components, one for ordinary representations and one for non-ordinary representations.

We are not quite done, because it is necessary to show that  $S_\emptyset$  is supported on both components. A simple argument shows that it is supported on the non-ordinary component (in brief: it is easy to see that there is a congruence to a cuspidal potentially crystalline representation, which after a further base change gives a non-ordinary crystalline representation), but in general to show that it is supported on the ordinary component requires the techniques of the third method discussed below.

The second method is that of [Kis09a], which applies only in the case that  $p$  splits completely in  $F$ , but again proves essentially complete results in this case. This method relies on the  $p$ -adic local correspondence for  $\mathrm{GL}_2(\mathbb{Q}_p)$  (which is the main reason for the restriction to the case that  $p$  splits completely in  $F$ ), as well as some more commutative algebra, and some results on the weight part of Serre’s conjecture (see below).

One disadvantage of both of these methods that they do not (at least at present) seem to generalise any further, and in particular they cannot be applied to  $n$ -dimensional representations when  $n > 2$ . The third method is that of [BLGGT10], which has the advantage of working for some  $n$ -dimensional potentially crystalline representations with arbitrary Hodge–Tate weights, but has the disadvantage that it is at present unclear how general a class of representations it applies to. Suppose that both  $\rho$  and  $\rho_0$  are unramified outside of  $p$ ; then our main argument shows that we can deduce the modularity of  $\rho$  from  $\rho_0$ , provided that for each place  $v|p$ ,  $\rho|_{G_{F_v}}$  and  $\rho_0|_{G_{F_v}}$  correspond to points on the same component of the local deformation ring. There is a definition of a “potentially diagonalizable” representation of  $G_{F_v}$ , which is one that (perhaps after base change) lies on the same component as a sum of characters. The basic idea is then that if the global representation  $\bar{\rho}$  happened to be induced from a character, then we could take  $\rho_0$  to be induced from a character, and if  $\rho|_{G_{F_v}}$  is potentially diagonalizable for

each  $v|p$ , we could deduce the modularity of  $\rho$  from that of  $\rho_0$  (which is known, as an instance of automorphic induction).

Of course, in general  $\bar{\rho}$  will not be induced from a character. However, in the general case, this problem can be circumvented by tensoring  $\bar{\rho}$  with a representation induced from a character, and using an idea of Michael Harris to “undo” this after proving a modularity lifting theorem. (Of course, this process turns a 2-dimensional representation into a 4-dimension representation, so it is necessary to develop a theory of higher dimensional automorphy lifting theorems in order to use this argument.) The eventual output is a modularity lifting theorem (in any dimension) in which it is necessary to assume that both  $\rho$  and  $\rho_0$  are potentially diagonalizable at all places dividing  $p$ , but where there are no assumptions on the local fields  $F_v$ , and no assumptions that the Hodge–Tate weights of  $\rho$  and  $\rho_0$  agree. (The reason that it is no longer necessary to assume that  $\rho, \rho_0$  have the same Hodge–Tate weights is that as part of the argument, both  $\rho, \rho_0$  are tensored with inductions of characters, and those inductions are not required to have the same Hodge–Tate weights.)

Unfortunately, it seems to be hard to establish potential diagonalizability in any generality, although it seems reasonable to expect that it holds for all potentially crystalline representations. It is easily checked to hold in both the ordinary and Fontaine–Laffaille cases, and this makes it a very useful notion in proving potential automorphy theorems for compatible systems of Galois representations; it turns out to be relatively easy to prove such theorems for ordinary representations, but unfortunately it is hard to prove that many Galois representations in a compatible system are ordinary (although it is conjectured that they are). On the other hand, all but finitely many of the representations are Fontaine–Laffaille, and potential diagonalizability gives a bridge from the ordinary to the Fontaine–Laffaille case.

This third argument gives one case in which it is not necessary to assume that  $\rho$  and  $\rho_0$  have the same Hodge–Tate weights. For many reasons, it would be desirable to remove this assumption whenever it is present in a modularity lifting theorem. This seems to be a hard problem in general, and is known as the weight part of Serre’s conjecture. It is now known in enough generality to remove it as an assumption from our main theorem, but the only known proof in this case makes use of ideas and methods from all three arguments above, together with the method of Khare–Wintenberger discussed below. It is plausible that a complete understanding of this problem within reach, but it seems to be bound up surprisingly tightly with the problem of proving general modularity lifting theorems where the Hodge–Tate weights of  $\rho, \rho_0$  are assumed to be equal.

**8.2. Further generalisations.** Other than the results discussed in the previous subsection, there are a number of obvious generalizations that one could hope to prove. One obvious step, already alluded to above, is to replace 2-dimensional representations with  $n$ -dimensional representations. This was originally done in [CHT08], [Tay08], and has been refined in a number of papers, most notably [Tho10], [BLGGT10]. It is necessary to assume that the Galois representations are essentially self-dual (that is, self-dual up to twist; note that this is automatic if

$n = 2$ ), but the argument then goes through in much the same way (albeit with a great deal more work!).

Another obvious way in which one could hope to relax the hypotheses in the theorem would be to allow  $F$  to be a more general number field. If  $F$  is not a CM field, then this seems to be out of reach at present, as there is not even a conjectural method for attaching Galois representations to automorphic representations. If  $F$  is a CM field, then the prospects are better; recently announced work of Harris–Lan–Taylor–Thorne constructs the required Galois representations (even for  $GL_n$ , without any self-duality hypothesis), and Calegari–Geraghty have suggested a strategy to adapt the modularity lifting machinery to this case, related to their work on weight one modular forms mentioned below. However, their strategy would require a generalisation of the methods of Harris–Lan–Taylor–Thorne to attach Galois representations to torsion classes in cohomology, so it seems likely that we are still some way off having general theorems here.

Another natural condition to relax would be the condition that the Hodge–Tate weights are distinct; for example, one could ask that they all be equal, and hope to prove Artin’s conjecture. For general  $n$ -dimensional representations, this appears to be completely beyond reach, but in the case  $n = 2$  the problems are more manageable. Since we do not know how to associate Galois representations to Maass forms, we stick with modular forms of partial weight 1; for simplicity, assume in fact that  $F = \mathbb{Q}$ , so that we want to prove a modularity lifting theorem for weight 1 modular forms.

The first problem that occurs when trying to adapt the arguments above is that weight 1 modular forms are not in the image of the Jacquet–Langlands transfer from the automorphic forms on any definite quaternion algebra, so it is necessary to work directly with modular forms. This in itself is not a fundamental obstruction, and indeed the original modularity lifting theorem of Taylor–Wiles was proved directly on the modular curve. However, the difference between weight 1 modular forms and higher weight modular forms is that the weight 1 forms are only seen in coherent cohomology, and they contribute to cohomology in degrees 0 and 1 (while higher weight cuspforms only contribute to cohomology in a single degree). This means that, for example, the analogue of Proposition 5.3 is false. (In particular, it is easy to find examples of mod  $p$  modular forms of weight 1 which do not lift to characteristic zero modular forms of weight 1.)

In addition, the corresponding local deformation ring has lower dimension than in the higher weight case, meaning that the numerical coincidence that  $\dim \mathcal{J}_\infty = \dim R_\infty$  no longer holds, and the commutative algebra arguments that we made above failed. Recently Calegari and Geraghty ([CG12]) found a way to avoid these problems, and prove a modularity lifting theorem for weight one modular forms. They do this by patching cohomology in degrees 0 and 1 simultaneously, effectively patching complexes of modules rather than modules. (There is also earlier work of Buzzard–Taylor, which has now been considerably generalised, which

proves a modularity lifting theorem more generally for weight 1 modular forms via one for Hida families and an analytic continuation argument.)

Finally, of course in the end we would like to be able to dispose of the hypothesis that  $\bar{\rho}$  is modular (that is, to dispose of  $\rho_0$ ). This is the problem of Serre's conjecture and its generalisations, and has only been settled in the case that  $F = \mathbb{Q}$  and  $n = 2$ . The proof in that case (by Khare–Wintenberger and Kisin) makes essential use of modularity lifting theorems and of the Khare–Wintenberger lifting method discussed below, in order to inductively reduce to the cases that  $p \leq 5$  and  $\bar{\rho}$  has very little ramification, when direct arguments using discriminant bounds can be made. The modularity lifting theorems discussed above make it plausible that the inductive steps could be generalised, but the base case of the induction seems specific to the case of  $\mathrm{GL}_2/\mathbb{Q}$ , and proving the result in greater generality is one of the biggest open problems in the field.

#### REFERENCES

- [BC] Olivier Brinon and Brian Conrad, *CMI summer school notes on  $p$ -adic Hodge theory*, available at <http://math.stanford.edu/~conrad/>.
- [BD12] Christophe Breuil and Fred Diamond, *Formes modulaires de Hilbert modulo  $p$  et valeurs d'extensions Galoisiennes*, 2012.
- [BLGG13] Tom Barnet-Lamb, Toby Gee, and David Geraghty, *Serre weights for rank two unitary groups*, *Mathematische Annalen* (to appear) (2013).
- [BLGGT10] Tom Barnet-Lamb, Toby Gee, David Geraghty, and Richard Taylor, *Potential automorphy and change of weight*, 2010.
- [BLGHT09] Tom Barnet-Lamb, David Geraghty, Michael Harris, and Richard Taylor, *A family of Calabi-Yau varieties and potential automorphy II*, to appear *P.R.I.M.S.*, available at <http://www.math.harvard.edu/~rtaylor>, 2009.
- [Boe] Gebhard Boeckle, *Deformations of Galois representations*, available at <http://www1.iwr.uni-heidelberg.de/groups/arith-geom/home/members/gebhard-boeckle/publications/>.
- [Buz12] Kevin Buzzard, *Potential modularity—a survey*, *Non-abelian fundamental groups and Iwasawa theory*, *London Math. Soc. Lecture Note Ser.*, vol. 393, Cambridge Univ. Press, Cambridge, 2012, pp. 188–211.
- [Car94] Henri Carayol, *Formes modulaires et représentations galoisiennes à valeurs dans un anneau local complet*,  *$p$ -adic monodromy and the Birch and Swinnerton-Dyer conjecture* (Boston, MA, 1991), *Contemp. Math.*, vol. 165, Amer. Math. Soc., Providence, RI, 1994, pp. 213–237.
- [CG12] Frank Calegari and David Geraghty, *Modularity lifting beyond the Taylor–Wiles method*, 2012.
- [Cho09] S.-H. Choi, *Local deformation lifting spaces of mod  $l$  Galois representations*, Ph.D. thesis, Harvard University, Cambridge, Massachusetts, 2009.
- [CHT08] Laurent Clozel, Michael Harris, and Richard Taylor, *Automorphy for some  $l$ -adic lifts of automorphic mod  $l$  Galois representations*, *Pub. Math. IHES* **108** (2008), 1–181.
- [Clo90] Laurent Clozel, *Motifs et formes automorphes: applications du principe de fonctorialité*, *Automorphic forms, Shimura varieties, and  $L$ -functions*, Vol. I (Ann Arbor, MI, 1988), *Perspect. Math.*, vol. 10, Academic Press, Boston, MA, 1990, pp. 77–159.
- [CR62] Charles W. Curtis and Irving Reiner, *Representation theory of finite groups and associative algebras*, *Pure and Applied Mathematics*, Vol. XI, Interscience Publishers, a division of John Wiley & Sons, New York-London, 1962.
- [DDT97] Henri Darmon, Fred Diamond, and Richard Taylor, *Fermat's last theorem*, *Elliptic curves, modular forms & Fermat's last theorem* (Hong Kong, 1993), *Int. Press*, Cambridge, MA, 1997, pp. 2–140.

- [Dia97] Fred Diamond, *The Taylor-Wiles construction and multiplicity one*, Invent. Math. **128** (1997), no. 2, 379–391.
- [Dic01] Mark Dickinson, *On the modularity of certain 2-adic Galois representations*, Duke Math. J. **109** (2001), no. 2, 319–382.
- [DT94] Fred Diamond and Richard Taylor, *Lifting modular mod  $l$  representations*, Duke Math. J. **74** (1994), no. 2, 253–269.
- [Far] Laurent Fargues, *Motives and automorphic forms: the (potentially) abelian case*, available at <http://www-irma.u-strasbg.fr/~fargues/Notes.html>.
- [FL82] Jean-Marc Fontaine and Guy Laffaille, *Construction de représentations  $p$ -adiques*, Ann. Sci. École Norm. Sup. (4) **15** (1982), no. 4, 547–608 (1983).
- [FM95] Jean-Marc Fontaine and Barry Mazur, *Geometric Galois representations*, Elliptic curves, modular forms, & Fermat’s last theorem (Hong Kong, 1993), Ser. Number Theory, I, Int. Press, Cambridge, MA, 1995, pp. 41–78.
- [Fon94] Jean-Marc Fontaine, *Le corps des périodes  $p$ -adiques*, Astérisque (1994), no. 223, 59–111, With an appendix by Pierre Colmez, Périodes  $p$ -adiques (Bures-sur-Yvette, 1988).
- [Kis08] Mark Kisin, *Potentially semi-stable deformation rings*, J. Amer. Math. Soc. **21** (2008), no. 2, 513–546.
- [Kis09a] ———, *The Fontaine-Mazur conjecture for  $GL_2$* , J. Amer. Math. Soc. **22** (2009), no. 3, 641–690.
- [Kis09b] ———, *Modularity of 2-adic Barsotti-Tate representations*, Invent. Math. **178** (2009), no. 3, 587–634.
- [Kis09c] ———, *Moduli of finite flat group schemes, and modularity*, Ann. of Math. (2) **170** (2009), no. 3, 1085–1180.
- [KW09] Chandrashekhara Khare and Jean-Pierre Wintenberger, *Serre’s modularity conjecture. II*, Invent. Math. **178** (2009), no. 3, 505–586.
- [Maz89] B. Mazur, *Deforming Galois representations*, Galois groups over  $\mathbf{Q}$  (Berkeley, CA, 1987), Math. Sci. Res. Inst. Publ., vol. 16, Springer, New York, 1989, pp. 385–437.
- [Maz97] Barry Mazur, *An introduction to the deformation theory of Galois representations*, Modular forms and Fermat’s last theorem (Boston, MA, 1995), Springer, New York, 1997, pp. 243–311.
- [Mil06] J. S. Milne, *Arithmetic duality theorems*, second ed., BookSurge, LLC, Charleston, SC, 2006.
- [Pil] Vincent Pilloni, *The study of 2-dimensional  $p$ -adic Galois deformations in the  $l$  not  $p$  case*, available at <http://perso.ens-lyon.fr/vincent.pilloni/>.
- [Ram93] Ravi Ramakrishna, *On a variation of Mazur’s deformation functor*, Compositio Math. **87** (1993), no. 3, 269–286.
- [Ser79] Jean-Pierre Serre, *Local fields*, Graduate Texts in Mathematics, vol. 67, Springer-Verlag, New York, 1979, Translated from the French by Marvin Jay Greenberg.
- [Sho13] Jack Shotton, in preparation, 2013.
- [SW99] C. M. Skinner and A. J. Wiles, *Residually reducible representations and modular forms*, Inst. Hautes Études Sci. Publ. Math. (1999), no. 89, 5–126 (2000).
- [SW01] C. M. Skinner and Andrew J. Wiles, *Nearly ordinary deformations of irreducible residual representations*, Ann. Fac. Sci. Toulouse Math. (6) **10** (2001), no. 1, 185–215.
- [Tay03] Richard Taylor, *On icosahedral Artin representations. II*, Amer. J. Math. **125** (2003), no. 3, 549–566.
- [Tay08] ———, *Automorphy for some  $l$ -adic lifts of automorphic mod  $l$  Galois representations. II*, Pub. Math. IHES **108** (2008), 183–239.
- [Tho10] Jack Thorne, *On the automorphy of  $l$ -adic Galois representations with small residual image*, 2010.

*E-mail address:* toby.gee@imperial.ac.uk

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON